A Direct and Simplistic Bromination of Commercially Important Organic Compounds in Aqueous Media by Eco-friendly $\text{AlBr}_3\text{-Br}_2$ Reagent System

Sushil Kumar Sharma^{1,*}, D. D Agarwal²

¹Department of Chemistry, Shri Jagdishprasad Jhabarmal Tibrewala University, Rajasthan, India

²Shri Jagdishprasad Jhabarmal Tibrewala University, Rajasthan, India

Copyright © 2015 Horizon Research Publishing All rights reserved.

Abstract A facile, simplistic, highly efficient, environmentally safe, regioselective, controllable and economical method for the bromination of organic compounds using aqueous $\text{AlBr}_3\text{-Br}_2$ reagent system.

Keywords Halogenation, Oxidative-Bromination, Molecular Bromine, Aqueous Medium, New Reagent System

1. Introduction

To date, available research on bromination of organic aromatic compounds in aqueous medium is very few and far between, except only few examples. Ganchegui et al. described the oxybromination of phenol ($\text{C}_6\text{H}_5\text{OH}$) and aniline $\text{C}_6\text{H}_5\text{NH}_2$ subsidiary products using $\text{NaBr-H}_2\text{O}_2$ in $\text{H}_2\text{O/scCO}_2$ (water/supercritical carbon dioxide) biphasic system, howbeit low conversion rate, high temperature (40°C), longer duration (from 4 to 20 h) and use of surplus amount of reagent (substrate : $\text{NaBr} : \text{H}_2\text{O}$; 1 : 3 : 3) are some of the concomitant inadequacies. Furthermore, experiments using dense gases must only be conducted in appropriate equipment and under apposite safety precautions. In recent times, Padgorsek et al. defined an interesting oxidative bromination of triggered arenes using $\text{H}_2\text{O}_2\text{-HBr}$ "on water" system is advantageous since water is the only by-product; though, a very high reaction time (from 8 hr to 28 hr) and the several threats associated with H_2O_2 make the process of limited industrial utility. The highly lethal 48 per cent aqueous HBr reacts aggressively with many metals with the generation of extremely flammable hydrogen gas, which may burst, which further limits its usage. Other described—on water || and —in water || brominating system are limited only to the synthesis of bromohydrins, α -bromoketones, iodination and benzylic bromination.

Factually hundreds of Br_2 -based and oxidative

bromination reagents have been described for the bromination of organic compounds, some latest reports are given in Table no. 5. For Bromine (Br_2) there are several other reagent systems that have been established as a substitute, but not restricted to, NBS/[Bmim]Br , $\text{ZrBr}_4/\text{diazene}$, $[\text{k.18-crown-6}]\text{Br}_3$, $[\text{BMPy}]\text{Br}_3$ $[\text{Hmim}]\text{Br}_3$, $[\text{Bmim}]\text{Br}_3$, 53f poly(4-vinylpyridinium tribromide), DABCO-bromine, pentapyridinium tribromide, ethylene bis(N-methylimidazolium) ditribromide. The bromination of polymer latex has been carried out using bromine emulsion which is prepared by liquefying bromine (Br_2) into an aq. solution of wetting agent/surfactant.

On the other hand, nothing like such reagent commercialized till date, because of its costly nature, very bad retrieval and recycling of used reagent/agent, waste treatment for large quantities of HBr -byproduct and poor stability of reagents and storage challenges in long period; therefore they are suitable for laboratory-scale requirements only with narrow application range. In the Previous section of the chapter, we have described aq. $\text{AlBr}_3\text{-Br}_2$ system in acetonitrile (MeCN) as a highly effective brominating agent which provided a rapid bromination of industrially important compounds in outstanding yields and pureness. The method is advantageous due to the cost-effective nature of the brominating reagent and high pureness of the wanted products, whereas the usage of organic solvents acetonitrile (MeCN) and the further utility. Consequently, to make the system more industrially-oriented and environmentally-approachable, we tried the bromination of liquid substrates using aq. $\text{AlBr}_3\text{-Br}_2$ can diffuse through the dews of aniline leading to bromination. During the reaction process the bromine (Br_2) color vanished instantly resulting the instantaneous synthesis of 2,4,6-tribromoaniline in 96 per cent yield (HPLC pureness 99.7 per cent) within 15 min of reaction time (Table no. 7, entry 19). This result heartened us to brominate the solid substrate (4-nitroaniline) under the similar conditions. As we noticed an instantaneous loss of reddish-brown hue in the round bottom flask and whole Br_2

get used within 2-3 minutes of magnificent signifying a prompt interface between the Br_2 and 4-nitroaniline has occurred in the aqueous system. Yellow color crystalline powder in 98 per cent yield of 2,6-dibromo-4-nitroaniline (HPLC purity of 99.08 per cent) was achieved (Table no. 7, entry 16). Cheered by the above resulted outcomes, the bromination of industrially-important compounds using aqueous Al Br_3 - Br_2 solution under aqueous conditions at room temperature without using any organic solvent before and after the completion of reaction, the reaction have successfully been performed

While bromination was carried out in water using aq. Al Br_3 system, it offers the following benefits:

(1) absence retrieval step and loss of solvent during the recycling procedure that makes the present reaction more appropriate, (2) no addition of water is obligatory after completion of reaction as the brominated product too is insoluble in water in case of acetonitrile solvent, 15 ml water has to be added for 10 ml acetonitrile (MeCN) solvent to separate the precipitated product. This upsurges the overall volume of filtrate thereby upturn the cost of recovery of Al Br_3 from the aq. medium at the completion of reaction, (3) the reaction is functionally-easy and friendly which don't requires heating or cooling, (4) the brominated compound and Al Br_3 recuperated at the end of reaction are in unadulterated form as there is no organic solvent in the reaction mixture for their adulteration and also the only HBr byproduct present in the aqueous filtrate has to be transformed to Al Br_3 (5) the present method leads to zero waste discharge to the surroundings, since from the three components of the reagent, i.e., Al Br_3 , Br_2 and water; one atom of Br_2 was shifted to organic substrate and the other half-forming HBr is again employed back to Al Br_3 by its neutralization with $\text{Ca}(\text{OH})_2$.

The Al Br_3 of the fresh reaction and additional Al Br_3 produced from HBr nullification was recovered as a crystalline solid by concentrating the aqueous filtrate (14 g Al Br_3 in 20 ml H_2O similar to collected works procedure). The recovered Al Br_3 has possible industrial value in different sectors i.e; Pharmaceuticals, preservers, and fire retardants etc. have also been employed effectively to rejuvenate the brominating reagent for bromination in consequent batches without any substantial loss of reactivity. Henceforward, at the end of the reaction we have nonentity to dispose to the environment which is justified in view of green chemistry.

Bromination of substituted aromatics compounds using aqueous Al Br_3 - Br_2 system

2. Objective

Clean and green reaction methods that do not use dangerous organic solvents are reinvigorated and are in excessive demand now. A modern report on "green solvents" determined that the use of solvents like dioxane, acetonitrile, 1-butanol, propanol, acids, acetone, ethyl, formaldehyde and

THF are not suggested from an environmental viewpoint. Water, on the other hand, is an auspicious solvent because it is readily available, non-flammable, non-toxic and can give the easy split-up of catalysts or reagents from many aromatic products. To date, water is rarely used in accessible research on bromination of aromatic compounds, which is very limited except few examples. Factually hundreds of Br_2 -based and oxidative bromination reagents have been described for the bromination of organic compounds. On the other hand, no such commercial reagent is available till date, due to their costly nature, poor retrieval and recycling of used reagent, waste treatment for large quantities of HBr -byproduct and poor stability of reagents and storage challenge in long periods; therefore they are suitable for laboratory-scale requirements only with narrow application range. The usage of aqueous AlBr_3 - Br_2 system as the most inexpensive brominating agent and H_2O as a reaction media exemplifies the most logical and reasonable choice of brominating reagent to fulfill the need of inexpensive, cleaner and most effective system. In simple words, H_2O fulfills all the requirements in direct synthesis of commercially- important brominated aromatic compounds. In current study, we are presenting, for the first time, a simple and direct bromination of $\text{C}_6\text{H}_5\text{OH}$ and $\text{C}_6\text{H}_5\text{NH}_2$ subsidiary products with strong electron-withdrawing groups (EWG) such as carboxylic ($-\text{COOH}$), nitro ($-\text{NO}_2$) and formyl ($-\text{CHO}$) as examples of pharmaceutical reaction intermediates under absolutely aqueous conditions.

3. Experimental

i) Reagents and analytics

Reagents and initial material were procured from across, Aldrich and Merck and were used as usual. Only fine powdered form substrates were used during complete reaction process, granulated and scaly substrates were grinded and transformed into fine powder prior to reactions to improve the dissolution factor. Twice as distilled water was used during the complete study, Water model no. 2695 instrument with PDA detector was used for HPLC analysis purpose, column C18 (250mm $\times$ 4.6 $\times$ 5 μ), solvent scheme of 70 per cent CH_3OH + 30 per cent H_2O , and 1 mL/minute flow rate. HPLC purity is represented by area per cent. Bruker Avance II 400 NMR spectrometer was used for NMR studies and chemical shifts were described by δ ppm; spectra were recorded in DMSO and CDCl_3 , ^1H NMR (comparative to TMS referenced as 0.00 ppm) and ^{13}C NMR (comparative to DMSO referenced as 39.50 ppm). GCMS studies were performed by using "Agilent GC-5893" with chemstation software; HP5-MS-column, with specification 30m $\times$ 0.25mm $\times$ 0.25 μ ; constant flow of- 2 ml/minutes; mass-director; mass range- 14 to 650 amu; detector temp- 290 $^\circ\text{C}$; injector temp- 280 $^\circ\text{C}$; injected sample volume- 1 microliter of 5 per cent solution in methanol. Mass spectroscopical (MS) data were documented on "Micromass Quattro Micro API triple quadrupole MS" which was

equipped with a ordinary APCI ion source. Infrared spectra studies were carried out on a Shimadzu Prestige 21 FT-IR Instrument (KBr, 3400-430 cm^{-1}). Isolated compounds were acknowledged on the basis of physical determination and spectroscopic data (13C NMR, 1H NMR, Infrared and Mass Spectroscopy).

ii) A typical synthesis path of 2, 6-dibromo-4-nitroaniline (16)

A solution was prepared using AlBr_3 (3.95 g, 20 mmol) in aqueous media (10 ml) and was added to Br_2 (3.25 g, 20 mmol), the resultant mixture was stirred continuously at room temperature till it give clear dark pale yellow solution. Immediately this solution was added to a fine powdered form of 4-nitroaniline (1.3733 g, 10 mmol) booked in a round-bottom flask (100 ml capacity) equipped with a magnetic stirrer using a pressure-equalizing funnel within 2 -3 minutes of time. The bromine color vanished immediately and 2,6-dibromo-4-nitroaniline was achieved (yellowish thick precipitates) within 20 minutes (Monitored by Thin Layer Chromatography) at room temperature. The filtration of precipitated mass was done by vacuum filtration method using Bunchner funnel, and then washed-off with $\text{Na}_2\text{S}_2\text{O}_3$ solution and dehydrated in oven at 100°C. The filtrate was saved for the Later runs. The product was achieved as yellow powder (2.90 g, 98 per cent yield, and 99.03 per cent HPLC area pureness). Melting point: 206°C (literature,²³ b 206-208°C). 1H NMR (400 MHz, DMSO): δ value 8.25 (s, 2H, ArH), 6.85 (s, 2H, NH_2); 13C NMR (100 MHz, DMSO): 144.36, 135.69, 122.49, 109.44; IR (KBr): 3450, 3366, 3184, 2798, 2659, 2363, 1642, 1515, 1488, 1348, 1298, 1270, 1154, 953, 879, 816, 747, 669, 584, 543, 464 cm^{-1} ; MS, Atmospheric Pressure Chemical Ionization (APCI) m/z (mass-to-charge ratio) called for $\text{C}_6\text{H}_4\text{Br}_2\text{N}_2\text{O}_2$:295.9; found 295.2.

iii) Procedure for recycling of Hydrogen Bromide as AlBr_3 and reformation of brominating reagent (Recycle 1)

The neutralization of filtrate originated from above reaction done by $\text{Al}(\text{OH})_3$ (0.7401 g, 10 mmol) to convert hydrogen bromide into AlBr_3 . Consequently, (3.23 g, 20 mmol) Br_2 was added to the aq. solution containing recycled aluminium tribromide solution was then added fine powder of 4-nitroaniline (1.3813 g, 10 mmol) quickly within 5 minutes taken in a round-bottom flask (100 ml) equipped with a pressure-equalizing funnel and magnetic stirring bar. The thick yellowish precipitates of 2, 6-dibromo-4-nitroaniline were attained within 20 minutes at room temperature immediately after adding Br_2 color disappeared. By using vacuum filtration the end product was separated from the mother liquor (solution left over after crystallization) and then washed-off with sodium thiosulfate ($\text{Na}_2\text{S}_2\text{O}_3$) solution (10 per cent, 10 ml $\times$ 3) and dried out in vacuum drying oven at 100°C. A high purity (99 per cent) 2, 6-dibromo-4-nitroaniline yellow powder was achieved in 2.82 g (98.08 per cent) yield with melting point 206°C. The distinguishing data was documented for the isolated product and found similar as given in the above distinctive method.

The hydrogen bromide evolved was again nullified and the aqueous solution was uninterruptedly recycled in the next run with an additional amount of Br_2 .

iv) The Route for Recycle/Reutilize 2,3 and 4

Analogous to the above typical method of Reutilize 1, Br_2 (3.2 g, 20 mmol) was added to the aq. solution containing recycled aluminium tribromide (AlBr_3) achieved after the separation of 2, 6-dibromo-4-nitroaniline ($\text{Br}_2\text{C}_6\text{H}_2(\text{NO}_2)\text{NH}_2$) and the reaction proceeded in a similar reaction pattern with 4- nitroaniline, $\text{C}_6\text{H}_5\text{N}_2\text{O}_2$ (1.3726 g, 10 mmol) for every reaction cycle.

4. Result & Discussion

i) Reaction conditions screening

The dibromination of para-substituted substrate decreases the possible complex mixture of mono- and dibrominated species, thus in order to abridge the analytics, 4- nitroaniline 16 (4-NA) was castoff as a typical substrate in a first-screening in a first- screening for appropriate reaction conditions. To optimize the yield and pureness, the bromine (Br_2) and (AlBr_3) concentration effects on the end product yield and Mp of 2, 6-dibromo-4- nitroaniline (DBNA) were studied in the dibromination of 4-NA.

ii) The effect on yield and melting point of DBNA by mole ration of Br_2

The excellence of the end product is strongly reliant on the mole ratio of Br_2 /4-NA which also has been confirmed by recent studies. It also been initiated that the optimal yield (98 per cent) of 2, 6-dibromo-4- nitroaniline (DBNA) and the desired melting point of 206°C (literature, 206-208°C) are achieved at 2:1 mole ratio of Br_2 /4-NA in the bromination of 4-NA by aqueous AlBr_3 - Br_2 solution as a effective brominating agent. Further, if we increase the mole ratio of Br_2 /4- NA from 2.0 to 2.2, yield of end product becomes stagnant. If we reduce the mole ratio from 2.0 to 1.8, the end product yield drops 98-93 per cent with a reduction in melting point 198-200 °C, which is not actually as per the required specifications. Further reduction in mole ratio of Br_2 /4- NA from 1.8 to 1.65, decrease the yield percentage of under-brominated product from 93-88 per cent, which melts at 160-170°C. The monobrominated 4-nitroaniline were achieved that melt at 102°C and 100-101°C, correspondingly (melting point of 2-bromo-4-nitroaniline is 104°C), at the mole ratio of 1.5 and 1.25.

iii) Influence of AlBr_3 - Br_2 mole ratio on yield and melting point of 2, 6-dibromo-4- nitroaniline (DBNA)

Available literature shows an identical pattern in the bromination of 4-NA using aqueous AlBr_3 solution as brominating agent because of increase in mole ratio of AlBr_3 /4-NA from 0.25 -2.0; at mole ratio 2:1 ideal yield and preferred melting point are achieved. With increase in mole ratio of AlBr_3 /4-NA from 0.25 to 2.0, melting point does not change significantly but the yield of the product upturns

from 91 to 98 per cent. The function of aluminium tribromide (AlBr_3) was defensible by performing a blank reaction for 1 h at 25°C using molecular bromine as a brominating agent which resulted in a composite fusion of under-brominated species that melt from 160 to 190°C . Thus, the result drawn in to conclusion that the optimal mole ratio 1:2:2 of 4-NA to AlBr_3 to Br_2 was establish to be best for the dibromination of 4-NA.

iv) Stirring

For mixing the hydrophobic aromatic substrate with the aqueous inorganic medium and for preventing local high concentrations of active bromo species, as the reaction was prompt, a proficient/effective stirring (750 rpm or higher) was required. The common applicability of this method was recognized when bromination of structurally-different activated aromatic compounds substituted with electron-withdrawing groups were analyzed (Table 2). It is the revolution of the present bromination reagent (aq. $\text{AlBr}_3\text{-Br}_2$ solution) over a newly reported brominating system ($\text{NBS}/[\text{Bmim}]\text{Br}$ or dioxane) that $\text{C}_6\text{H}_5\text{OH}$ and $\text{C}_6\text{H}_5\text{NH}_2$ byproducts with strong electron-withdrawing groups (EWG) such as $-\text{COOH}$, $-\text{NO}_2$ and $-\text{CHO}$ were promptly brominated regioselectively in outstanding yields and pureness. The same reported system, however, yields a combination mix of mono- and dibromo products. In the present system, salicylic acid 1 and 4-hydroxybenzoic acid 2 were transformed with quantitative conversion resulting in clean synthesis of 3,5- dibromosalicylic acid (92 per cent yield; 96.2 per cent HPLC purity) and 3,5-dibromo-4-hydroxybenzoic acid (94 per cent yield; 99.2 per cent HPLC purity) after 20 min and 15 min, respectively. Pharmaceutically-important aromatic aldehydes (Table 3, entries 3, 4, 5) were also instantaneously synthesized using aq. $\text{AlBr}_3\text{-Br}_2$ system in excellent yields. Under these conditions, acetaniline 6 and benzaniline 7 were

regioselectively transformed to their mono-brominated products in very good yields. The regioselectivity witnessed in these reactions matches to that predictable for an electrophilic bromination path modulated by steric effects. We observed that oxine (8) and sulfanilamide (9) could also be efficiently brominated affording pharmaceutically-important 5, 7-dibromooxine (96 per cent yield; 99.67 per cent HPLC pureness) and 3,5-dibromosulfanilamide (94 per cent yield; 96.6per cent HPLC pureness) within 15 minutes. Another antibacterial compound, 2, 4-dibromo-6- nitrophenol was isolated in excellent yield within 20 minutes from 2-nitrophenol (Table no. 6, entry 10). With the poor reactivity 4-nitrophenol (11) shows outstanding reactivity resulting 2,6-dibromo-4-nitrophenol (94 per cent yield; 97.5 per cent HPLC pureness) within 30 minutes using two equivalent of reagent. Bromination of the same substrate with $\text{H}_2\text{O}_2\text{-HBr}/\text{on water}$, gave 2, 6-dibromo-4-nitrophenol over a period of 24 hr. The advantages of the present system is an regioselective, effective, facillie, and fast bromination of deactivated anilines (Table no. 6, entries 12 to 16) in outstanding end product yields at 25°C upon simple admixing with aq. $\text{CaBr}_2\text{-Br}_2$ solution which is slightly challenging by other methodologies. 1-Bromo-2- naphthol 17 (92 per cent yield; 96.2 per cent HPLC pureness) was promptly obtained under undistinguishable reaction conditions in excellent yield within 10 minutes of reaction time while for 1,6- dibromo-2-naphthol 18 (97 per cent yield; 96.5 per cent HPLC pureness), within 25 minutes and 2 equivalents of aqueous $\text{AlBr}_3\text{-Br}_2$ solution were necessary. Which represents the position of the electrophilic attack and the number of inflowing Br_2 atoms during reaction process can be structured by controlling the ratio of substrate: aqueous $\text{AlBr}_3\text{-Br}_2$ solution, i.e., 1/1 mono-, 1/2 for di-, and 1/3 for tribromination of aromatic compounds.

Table 1. A comparison of the results of present system (aq. $\text{AlBr}_3\text{-Br}_2$ solution) with the literature precedents of some recently published brominating systems in the synthesis of pharmaceutical intermediates

Entry	Substrate	Product	aq $\text{AlBr}_3\text{-Br}_2$ (present system) ^a	Recently published brominating systems
1.			92per cent, 15 min, 25°C(99.2per cent) ^b	$\text{NH}_4\text{Br}/\text{H}_2\text{O}_2/\text{AcOH}$ 73per cent, 4h, rt $\text{HBr}/\text{H}_2\text{O}_2/\text{MeOH}$ 99per cent, 12h, 0°C to rt, $\text{NBS}/\text{SO}_3\text{H}$ -functionalized silica 91per cent, 3h, rt.
2.			96per cent, 15 min, 25°C(99.9per cent) ^b	$\text{LiBr}/\text{CAN}/\text{MeCN}$ 99per cent, 1h, rt, $\text{KBr}/\text{BTPPMS}/\text{MeCN}$ 90per cent, 4h, rt, $\text{TEATB}/\text{H}_2\text{O-DMF}$ 65per cent, 1h, rt, $\text{KBr}/\text{H}_2\text{O}_2/\text{CuPcF}_{16}\text{-APSG}/\text{AcOH}$ 87per cent, 2.2h, 60 °C
3.			93per cent, 20 min, 25°C(96.6per cent) ^b	$\text{NBS}/\text{NH}_4\text{OAc}/\text{MeCN}$ 89per cent, 1h, rt $\text{NBS}/\text{MeCN}/\text{hv}$ 88per cent, 30 min, 32°C $\text{NBS}/\text{SO}_3\text{-functionalized silica}$ 73per cent, 3h, rt.
4.			93per cent, 15 min, 25°C(96.2per cent) ^b	Hexaminethylene tetraminebromine/ CH_2Cl_2 30per cent, 5 min, rt
5.			96per cent, 15 min, 25°C(99.67per cent) ^b	$\text{HBr}/\text{iso-amyl nitrite}/\text{CH}_2\text{Cl}_2$ 100per cent, 16h, rt
6.			94per cent, 15 min, 25°C(97.5per cent) ^b	$\text{H}_2\text{O}_2\text{-HBr}/\text{'on water'}$ 95per cent, 24h, rt
7.			96per cent, 15 min, 25°C(99.7per cent) ^b	$\text{HBr}/\text{H}_2\text{O}_2/\text{MeOH}$ 96per cent, 8h, 0°C to rt $\text{TEATB}/\text{H}_2\text{O-DMF}$ 63per cent, 1.5h, rt $\text{TEATB}/50\text{per cent aq DMF}$ 65per cent, 15 min, rt
8.			97per cent, 25 min, 25°C(96.5per cent) ^b	$\text{HBr}/\text{H}_2\text{O}_2/\text{MeOH}$ 84per cent, 20h, rt
9.			92per cent, 10 min, 25°C(96.23per cent) ^b	$\text{HBr}/\text{H}_2\text{O}_2/\text{MeOH}$ 93per cent, 15h, rt $\text{KBr}/\text{H}_2\text{O}_2/\text{AcOH}/\text{Zeolite}$ 45per cent, 5h, rt $\text{KBr}/\text{BTPPMS}/\text{MeCN}$ 92per cent, 2h, rt $\text{KBr}/\text{H}_2\text{O}_2/\text{CuPcF}_{16}\text{-APSG}/\text{AcOH}$ 75per cent, 2h, 60°C $\text{H}_2\text{O}_2\text{-HBr}/\text{'on water'}$ 90per cent, 8h, rt
10.			98per cent, 20 min, 25°C(99.03per cent) ^b	$\text{HBr}/\text{H}_2\text{O}_2/\text{MeOH}$ 95per cent, 10h, 0°C to rt $\text{H}_2\text{O}_2\text{-HBr}/\text{'on water'}$ 95per cent, 28h, rt

The use of aqueous $\text{AlBr}_3\text{-Br}_2$ system as the most economical brominating agent and H_2O as a solvent/reaction medium denotes the most logical choice as a brominating reagent to fulfill the need of inexpensive, cleaner and most effective system for the instantaneous synthesis of commercially-significant brominated compounds. A comparative study of the brominating capability of the aqueous $\text{AlBr}_3\text{-Br}_2$ system with the recently circulated methods is testified in Table 5 which clearly indicates the benefits of the present system over obtainable methods. In this present study, we are presenting, an effective reagent system for electron-withdrawing groups (EWG) like; $-\text{COOH}$, NO_2 and $-\text{CHO}$ as examples of pharmaceutical reaction intermediates under purely aqueous conditions.

Table 2. Bromination by using aqueous $\text{AlBr}_3\text{-Br}_2$ system for several aromatic compounds

Entry	Substrate	Product	Solvent system				Mp/°C (lit.)	Application
			Water ^a		Acetonitrile ^b			
			Time /min	Yield ^c (per cent)	Time /min	Yield ^c (per cent)		
1.			20	96	25	97	103-104 (105-107)	Antibacterial, antifungal
2.			15	93	10	96	83-84 (84)	Antibacterial, antifungal and anthelmintic
3.			18	95	08	96	120 (120-121)	Intermediate for agrochemical-s and pharmaceuticals
4.			20	93	20	94	104(102-104)	Intermediate for dyestuffs
5.			20	98	10	98	206(206-208)	A potent antifungal, in the preparation of diazonium
6.			18	91	15	90	108(110-113)	Fine organic and custom intermediate
7.			18	96	15	96	127(128-129)	Fine organic and custom intermediate
8.			15	96	15	95	101(101-103)	Precursor for substituted thiazoles used as fungicides
9.			30	94	28	95	145(145-147)	Pharmaceutic-al intermediate
10.			18	91	25	96	114(116-117)	Anthelmintic

11.			20	95	20	94	235(235-237)	Pharmaceutic-al intermediate
12.			15	96	15	95	200(198-200)	A potent antifungal antiamoebic
13.			18	97	20	95	200-202 (202)	Pharmaceutic-al intermediate
14.			15	95	12	98	166(165-169)	Analgesic, Antipyretic
15.			25	92	35	96	163(164-166)	In pharmaceutical flavor
16.			25	95	15	93	82(80-84)	Pharmaceutic-ally acceptable salt as inhibitor
17.			20	92	20	96	183(181-185)	Pharmaceutic-al intermediate
18.			20	91	20	90	226(226-229)	Bactericide when incorporated in to topical ointments
19.			20	95	---	---	270(271-274)	Medical intermediate

Underneath these reaction conditions, the deactivated substrate like nitrobenzene and benzoic acid did not show any transformation at all. Table no. 2 discloses that the yield and the reaction time for several brominated products are comparable in H₂O and acetonitrile (ACN) as solvents, correspondingly. The only limitation associated with the reaction using water as a solvent is that granular and scaly substrates have to be crushed and ground proceeding to reactions to convert them into fine powders. Since the reaction operates entirely in water and generates absolute zero discharge; it seems very effective valuable from environmental safety point of view to extend this system for other commercially-important compounds. Subsequent, a desirable greener approach to the present study is following by an environmental friendly recycling procedure. The nonexistence of organic waste and organic solvents in the reaction process supported simple separation procedure comprised of filtration of solid phase brominated compounds and the addition of calcium hydroxide powdered form

(Ca(OH)₂) to the filtrate for the neutralization of hydrogen bromide waste so that the bromine of hydrogen bromide thus fixed as Aluminum bromide. The aqueous filtrate was reclaimed in the next cycle and the reinforced brominating agent was then used to brominate the substrate successively in the next run.

Table 3. Reprocessing experiments for the dibromination of 4-nitroaniline

Recycle	Appearance	Yield ^b (per	Mp/ ^o C(lit.,	Purity ^c
Newbatch	Yellow Powder	98.2	206	99.20
Reprocess 1	Yellow Powder	98.0	206	98.98
Reprocess 2	Yellow Powder	97.6	204-206	98.57
Reprocess 3	Yellow Powder	97.3	205-206	98.26
Reprocess 4	Yellow Powder	97.4	204-206	98.84

^a Surroundings: 10mmol 4-NA, 20mmol Br₂ (4-NA and Br₂ moles charged in every reaction cycle) and 20mmol AlBr₃ (charged for fresh batch only) at room temperature (25°C) for 20 minutes (each cycle).

^b Inaccessible Yields

^c Purity determined by HPLC

By this methodology, very productively and effectively the dibromination of 4-nitroaniline has been carried out for 4 cycles after the fresh batch using 4-NA: Br_2 (1: 2) mole ratio in each run without adding of the fresh aluminium tribromide (AlBr_3) to give 2, 6-dibromo-4-nitroaniline within 20 minutes at room temperature. Reprocessing experiments (Table no. 7) shows that for at least 4 cycles followed by the fresh batch, there is no substantial loss of reactivity of the rejuvenated brominating reagent. When the concentration of aluminium tribromide (AlBr_3) upturns in the filtrate, then through vaporization process filtrate was concentrated which

causes precipitation of AlBr_3 as a crystalline solid and an added amount of AlBr_3 (7 mol) was recuperated after fresh batch + 4 cycles: resulted 1 mol of aluminium tribromide (AlBr_3) produced likewise in every cycle by the neutralization of hydrogen bromide by adding $\text{Al}(\text{OH})_3$. Therefore, starting with 1 mol of 4-NA wrt 2 mol of Aluminium tribromide in the fresh batch, At the end of reaction process we achieved 7 mol of AlBr_3 . Consequently, effectively the byproduct hydrogen bromide has been utilized, and at the end of reaction, process generates zero discharge of organic waste and effluent to the surroundings.

Characterization of Representative Brominated Compounds:

Figure 8. ^1H -NMR spectra of 3,5-dibromosalicylic acid (1)

Figure ¹H-NMR spectra of 4-bromoacetanilide (6)

Figure 9. IR spectra of 3,5-dibromosulfanilamide (9)

Line#:2 R.Time:14.850(Scan#:3556)

MassPeaks:164

RawMode:Single 14.850(3556) BasePeak:330(419935)

BG Mode:None Group 1 - Event 1

Figure 10. GC-MS of 3,5-dibromosulfanilamide (9)

Line#:2 R.Time:10.517(Scan#:2256)

MassPeaks:77

RawMode:Averaged 10.513-10.520(2255-2257) BasePeak:297(60688)

BG Mode:Calc. from Peak Group 1 - Event 1

Figure 11. GC-MS of 2,4-dibromo-6-nitrophenol (10)

Figure 12.LC/MS of 2,6-dibromo-4-nitroaniline ($\text{Br}_2\text{C}_6\text{H}_2(\text{NO}_2)\text{NH}_2$) (16)

Figure 13. ¹H and ¹³C-NMR of 2,6-dibromo-4-nitroaniline (16)

5. Conclusions

We have testified a rapid, mild, cost effective and facile method for the effective and selective synthesis of pharmaceutically-important brominated intermediates using aq. AlBr₃-Br₂ system as an instantaneous brominating agent under aqueous conditions. The green features of this practice include no use of organic solvent and an operative utilization of HBr byproduct which lead to zero organic waste and zero waste discharge to the environment, consequently applicable for large scale bromination.

The categorization data (1H NMR, Infrared and Mass Spectroscopy) achieved for various representative compounds are given below:

3,5-Dibromo-4-hydroxybenzoic acid (2): White fine crystalline powder; 1H NMR (400 MHz, Dimethyl Sulfoxide-DMSO): δ 8.02 (2H, ArH, s), 11 (1H, OH, s), 13 (1H, COOH, s); Infrared (KBr): 3450, 3081, 2976, 2652, 2530, 1798, 1690, 1588, 1482, 1420, 1404, 1340, 1305, 1275, 1205, 1169, 1137, 903, 893, 766, 738, 718, 671, 520, 470 cm⁻¹; MS m/z (mass-to-charge ratio) calcd. for C₇H₄Br₂O₃: 295.9, found 295.

5-Bromovanillin (5): Slightly-brown powder; IR (KBr): 3318, 3103, 3074, 3009,

2976, 2943, 2851, 2746, 1725, 1592, 1502, 1464, 1448, 1425, 1355, 1293, 1192,

1148, 1047, 973, 855, 796, 681, 602, 592, 538, 527 cm⁻¹; MS m/z (mass-to-charge ratio) calcd. For C₈H₇BrO₃: 231, found 235.

4-Bromoacetanilide (6): White crystals, 1H NMR (400 MHz, Dimethyl Sulfoxide-DMSO): δ 2.1 (s, 3H, CH₃), 7.41 (d, J = 2.96 Hz, 2H, Ar), 7.64 (d, J = 2.89 Hz, 2H, Ar), 9.71 (s, 1H, NH). IR (Potassium Bromide): 3288, 3261, 3157, 3128, 3045, 1684, 1656, 1594, 1576, 1544, 1478, 1388, 1312, 1280, 1246, 1002, 845, 824, 754, 678, 508 cm⁻¹. MS Atmospheric Pressure Chemical Ionization (APCI) m/z (mass-to-charge ratio) calcd. For C₈H₈BrNO: 216.07, found 216.

5,7-Dibromo-8-hydroxyquinoline (8): Light buff colored powder; 1H NMR (400 MHz, Trichlorodeuteromethane "CDCl₃"): δ value 7.446 (d, J = 0.514 Hz, 1H), 8.465 (ddd, J = 8.803, J = 1.836, J = 0.514 Hz, 1H), 7.223 (dd, J = 8.778, J = 4.681 Hz, 1H), 8.839 (dd, J = 4.687, J = 1.834 Hz, 1H). IR (KBr): 3087, 1742, 1576, 1489, 1464, 1387, 1374, 1266, 1212, 1156, 1043, 947, 884, 815, 782, 736, 672, 643, 614, 589, 559, 500 cm⁻¹. MS Atmospheric Pressure Chemical Ionization (APCI) m/z (mass-to-charge ratio) calcd. For C₉H₅Br₂NO: 302.95, found 302.2.

3,5-Dibromosulphanilamide (9): Shining-white crystalline powder; 1H NMR (400 MHz, Trichlorodeuteromethane "CDCl₃"): δ : 5.74 (2H, s, NH₂), 6.76 (s, 2H, SO₂NH₂), 7.92 (s, 2H, Ar); IR (KBr): 3443, 3341, 3238, 3076, 1614, 1545, 1464, 1395, 1317, 1151, 1114, 912, 883, 731, 580, 546, 496 cm⁻¹; MS Atmospheric Pressure Chemical Ionization (APCI) m/z (mass-to-charge ratio) calcd. for C₆H₆Br₂N₂O₂S: 330, found 330.

2,4,6-Tribromoaniline (19): Shining white fine needles;

1H NMR (400 MHz, Trichlorodeuteromethane "CDCl₃"): δ : 7.5 (2H, s, Ar), 4.54 (2H, br. S, NH₂). Infrared (KBr): 3436, 3288, 1468, 1392, 1274, 1238, 1057, 849, 736, 712, 666, 542, 478 cm⁻¹. MS Atmospheric Pressure Chemical Ionization (APCI) m/z (mass-to-charge ratio) calcd. for C₆H₄Br₃N: 329.816, found 328.8.

REFERENCES

- [1] De la Mare, P. B. Electrophilic Halogenation: Reaction Pathway Involving Attack by Electrophilic Halogens on Unsaturated Compounds, Cambridge University Press, Cambridge, UK, (1976), Chapter 5. (b) Taylor, R. Electrophilic Aromatic Substitution, Wiley, Chichester, UK, (1990).
- [2] a) Ganchegui, B.; Leitner, W. Green Chemistry (2007), 9, 26. (b) Podgorsek, A.; Stavber, S.; Zupan, M.; Iskra, Journal Tetrahendron (2009), 65, 4429.
- [3] Pingali, S. R. K.; Madhav, M.; Jursic, B.S. Tetrahendron Letters. (2010), 51, 1383.
- [4] a) Surine, W. R.; Majewski, T.E. Preparation of 3,5-dibromosalicylic acid. U.S. Patent 3,381,032, (1968). (b) Hai, T.T.; Nelson, D.J. Pharmaceutical grade 3,5-dibromosalicylic acid and method for synthesizing same. U.S. Patent 5,013,866, (1991).
- [5] (a) Bedekar, A. V.; Gadde, R.; Ghosh, P. K. Process for preparing 2,4,4,6-tetrabromo-2, 5-cyclohexadienone. U.S. Patent 6,838,582, (2005). (b) De La Mare, P. B. D. Electrophilic Halogenation: Reaction Pathway Involving Attack by Electrophilic Halogens on Unsaturated Compounds; Cambridge University Press: London.
- [6] (a) The Merck Index, 13th edition, An Encyclopedia of Chemicals, Drugs, and Biologicals, Merck and Co. Inc., Whitehouse Station, NJ, 2001. (b) Aldrich Handbook of Fine Chemical Company, Inc.: Wisconsin, USA, (1990).
- [7] Adimurthy, S.; Ramachandraiah, G.; Bedekar, A.V.; Ghosh, S.; Ranu, B.C.; and Ghosh, P.K.; (2006). Eco-friendly and versatile brominating reagent prepared from a liquid bromine precursor. Green Chemistry, 8, 916–922 | 917.
- [8] Zolfigol, M.A.; Chehardoli, G.; Salehzadesh, S.; Adams, H.; Ward, M. D. Tetrahendron Letters, (2007), 48, 7969.
- [9] Hosseinzadeh, R.; Tajbakhsh, M.; Mohadjerani, M.; Lasemi, Z. Synthesis. Communication, (2010), 40, 868.
- [10] Al-Zoubi, R.M.; Hall, D. Organic Letters (2010), 12, 2480
- [11] Beckmann, J.; Bolsinger, J.; Duthie, A.; and Finke, P.; (2013). Diarylhalotelluronium(IV) cations [(8-Me₂NC₁₀H₆)₂TeX]⁺ (X = Cl, Br, I) stabilized by intramolecularly coordinating N-donor substituents. Dalton Trans. 10.1039.
- [12] Chiappe, C.; Leandri, E.; Pieraccini, D. Chemistry Commun. (2004), 2536.
- [13] Beckmann, J.; Bolsinger, J.; Duthie, A.; and Finke, P.; (2013). Diarylhalotelluronium(IV) cations [(8-Me₂NC₁₀H₆)₂TeX]⁺ (X = Cl, Br, I) stabilized by intramolecularly coordinating N-donor substituents. Dalton Trans. 10.1039.

- [14] Benitez, F. J.; Acero, J.L.; Real, F.J.; Roldan, G. and Casas, F. (2011). Bromination of selected pharmaceuticals in water matrices. *Chemosphere*. 85: 1430-1437.
- [15] Borikar, S.P.; Daniel, T.; Paul, V. *Tetrahedron Letters*. (2009),50,1007.
- [16] Chinnagolla, R.K.; Pimparkar, S.; and Jeganmohan, M.; (2013). Ruthenium-catalyzed intramolecular selective halogenation of O-methylbenzohydroximoyl halides: a new route to halogenated aromatic nitriles. *Chem. Commun.*, 49, 3146-3148.
- [17] Ceska, G. W. Aqueous bromine emulsions. U.S. Patent 3,886,081, (1975). Chaudhuri, M. K.; Khan, A. T.; Patel, B. K. *Tetrahedron Letters*. (1998), 39, 8163.
- [18] Chiu, Y. (Denville, NJ); Cottrell, S.A. (Baton Rouge, LA); Tung, H. S. (Getzville, NY); 'Kopkalli, H.(Staten Island, NY) and Cerri, G.(Parsippany, NJ). (2011). Process for the Manufacture of Fluorinated Olefins. U.S. Class: 570/175; Serial No.: 402,372.
- [19] Chiappe, C.; Leandri; Pieraccini, D. *Chem. Commun.* (2004), 2536.
- [20] Chinnagolla, R.K.; Pimparkar, S.; and Jeganmohan, M.; (2013). Ruthenium- catalyzed intramolecular selective halogenation of O-methylbenzohydroximoyl halides: A new route to halogenated aromatic nitriles. *Chem. Communications*, 49, 3146—3148.
- [21] Sharma, S.; Agarwal, D. D.; (2014). NH₄Br-Br₂ catalyzed oxidative bromination of aromatic compounds. *Journal of Agriculture and Life Sciences*. Vol. 1, No. 1; 65-81.
- [22] Sharma, S.; Agarwal, D. D.; (2014). Oxidative Chlorination of aromatic compounds in aqueous media. *Journal of Agriculture and Life Sciences*. Vol. 1, No. 1; 146-164.
- [23] D Sharma, S.; Agarwal, D. D.; (2014). Molybdate catalyzed oxidative bromination of aromatic compounds using mineral acids and H₂O₂. *International Journal of scientific and research publications*. Vol. 4, Issue 6; 01-12.
- [24] Das, B.; Venkateswarlu, K.; Mahji, A.; Siddaiah, V.; Reddy, K. R. J. *Molecular Catalysis A: Chemical* (2007), 267, 30. Del Valle, C. J. Method of producing calcium bromide. U.S. Patent 4, 234, 556, 1980.
- [25] Deshmukh, A.P.; Pandiya, K. J.; Jadhav, V.K.; and Salunkhe, M.M.; (1998). Halogenation of Aromatic Compounds by using Sodium Perborate as an Oxidant. *Journal of Chemical Research (S)*, 828-829.
- [26] Do, H.Q.; Daugulis, O.; (2008). A Simple Base-Mediated Halogenation of Acidic sp² C-H Bonds under Noncryogenic Conditions. *Organic Letters*, Vol.11, No. 2, 421-423.
- [27] Eberlin, A.; Williams, D.L.H.; (2002). Halogenation of enol tautomers of 2- cyanoacetamide and malonamic acid. *Journal of Chem. Soc., Perkin Transactions*. 2, 1316-1319.
- [28] Edwards; Jeffrey, D.; Palermo; John. Analgesic and anti-inflammatory composition. U.S. Patent 7,510,732, (2009).
- [29] Fong, H.L.(Sugar Land, TX); Johnson, T.H.(Houston, TX) and Semple, T.C.(Friendswood,TX).(2009). Conversion of Alkylhalides in to Alcohol Alkoxylates. *Current U.S. Class*: 568/671. Serial No.: 887194.
- [30] Forsyth; Adam, B.; Pryor; Ernest, D.; McGarry; James, E.; Harney; Gerald, D. W. Compositions containing certain 2,4-halo-6-nitrophenols or derivatives thereof and methods for using same to eradicate internal parasites in warm- blooded animals. U.S. Patent 4,031,249, 1977.
- [31] Gavara, L.; Boisse, T.; Riogo, B.; Henichart, J. P. *Tetrahedron-SD* (2008), 64, 4999.III, D. W.; Eaves Jr., C. C.; Haynes Jr., J. F. *Annals of Emergency Medicine* 1989, 18, 778.
- [32] Gerhard, H.; Phillips; Gray, W; Pratt; John, K.; Srouji; Gabriel, H. Substituted thiazoles and their use as fungicides. U.S. Patent 5,045,554, (1991).
- [33] Gershon, H.; Parmegiani, R.; Godfrey, P.K. *Antimic. Agents Chemotherapy* (1972), 1, 373.
- [34] Gershon, H.; McNeil, M. W.; Parmegiani, R.; Godfrey, P. K. *Applied Microbiology* (1971), 22, 438.
- [35] Gnam, J. M.; Sheldon, R. A. *Tetrahedron Letters*. (2005), 46, 4465.
- [36] Hai, T. T.; Nelson, D. J. Pharmaceutical grade 3,5-dibromosalicylic acid and method for synthesizing same. U.S. Patent 5, 013, 866, (1991).
- [37] Hajipour, A.R.; Pourmousavi, S.A. and Ruoho, A.E. (2006). 1-Benzyl 4-aza-1-azoniabicyclo[2.2.2]Octanetribromide as a Regenerable and Useful Reagent for Bromination of Phenols under Mild Conditions. *Ind. Journal of Chem.* 45B (Mar): 796-800.