Skewable Matrices over $\wedge^2 V$ Applied to Locally Metric Connections

Mihail Cocos, Kent Kidman*

Department of Mathematics, Weber State University, USA

Copyright ©2017 by authors, all rights reserved. Authors agree that this article remains permanently open access under the terms of the Creative Commons Attribution License 4.0 International License

Abstract A necessary condition for a connection in a vector bundle to be locally metric is for its curvature matrix, which consists of 2 forms, to be skew symmetric with respect to some local frame. In this paper we give a simple algorithm that can be used to decide when a matrix of 2 forms is equivalent to a skew symmetric matrix. We apply this algorithm to verify whether a full rank curvature connection is locally metric.

Keywords Simultaneously Skew Matrices, Linear Connections, Metric Compatible Connections, Full Rank Curvature

1 Introduction

The problem of finding metrics compatible with a connection in a vector bundle has been addressed by some mathematical physicists and mathematicians. See for example [2],[3],[4],[5].

This is a problem motivated by gauge theory in physics and also interesting as a pure mathematical question. We will now briefly describe the differential geometry problem and abstract from it the algebra problem our paper is concerned with.

Let E be a vector bundle of rank m above the smooth n dimensional manifold M , and D be a connection in E . It is a standard fact in differential geometry that if the connection D is locally metric (see for example [1] pages 249-256) then there exists a local frame

$$\sigma = (\sigma_1, \sigma_2, \dots, \sigma_m)$$

in E such that the curvature matrix of D with respect to σ is skew symmetric. It has been shown in [2] that the skew symmetry of the curvature matrix with respect to some frame is, in the two dimensional case, also sufficient for the local metrizable of the connection. The curvature matrix Ω consists of differential 2 forms defined on M and hence at a point $p \in M$ the curvature has elements of $\wedge^2 T_p M = T_p M \wedge T_p M$ as its entries. The problem then is to be able to find out if there is a frame in which the curvature matrix is skew symmetric. Here we have to remind the reader that if $\sigma' = (\sigma'_1, \sigma'_2, \dots, \sigma'_m)$

is a different local frame of the bundle E then the curvature matrix with respect to this new frame is related to the curvature matrix Ω by the equation

$$\Omega' = U^{-1}\Omega U \quad (1.1)$$

where U is the matrix with scalar entries defined by

$$\sigma' = \sigma U. \quad (1.2)$$

The equation (1.1) is essential since in the case of a connection D one can hardly expect its curvature matrix to be skew symmetric with respect to an arbitrary frame σ' . Therefore starting with an arbitrary local frame σ' we would like to be able find the scalar matrix U as in (1.2) that will generate our desired frame σ . It is therefore necessary for us to address the pure algebraic question of whether a given matrix Ω' with entries 2 forms is equivalent as in (1.1) with a skew symmetric matrix Ω . Next section will clarify the algebra problem.

2 The matrix problem

Let K be an n -dimensional real vector space and $e = (e_1, e_2, \dots, e_n)$ a frame in K , and let $\wedge^2 K = K \wedge K$ denote the vector space of wedge products of vectors in K .

Definition 2.1. An $m \times m$ matrix Ω with entries in $\wedge^2 K$ is **skewable** if there is a non-singular $m \times m$ real matrix U such that $(U^{-1}\Omega U)^T = -U^{-1}\Omega U$, i.e. if $U^{-1}\Omega U$ is skew-symmetric.

Definition 2.2. Let $S = \{S_1, S_2, \dots, S_r\}$ be a set of $m \times m$ real matrices. Then S is simultaneously skewable if there exist a nonsingular real $m \times m$ matrix U such that for each $1 \leq k \leq r$, $U^{-1}S_k U$ is skew symmetric.

Lemma 2.3. Let $S = \{S_1, S_2, \dots, S_r\}$ be a set of $m \times m$ real matrices. Suppose

$$\Omega = w_1 S_1 + w_2 S_2 + \dots + w_r S_r,$$

with $w_k \in \wedge^2 K$ and for which the set $\{w_1, w_2, \dots, w_r\}$ is linearly independent. Then Ω is skewable if and only if S is simultaneously skewable.

Proof. Suppose S is simultaneously skewable, and that $U^{-1}S_k U$ is skew symmetric for $1 \leq k \leq r$, for a nonsingular real U . Let $\Omega = w_1 S_1 + w_2 S_2 + \dots + w_r S_r$, $w_k \in \wedge^2 K$, where the set $\{w_1, w_2, \dots, w_r\}$ is linearly independent. Consider

$$U^{-1}\Omega U = U^{-1}(\sum_{k=1}^r w_k S_k)U = \sum_{k=1}^r w_k U^{-1}S_k U.$$

Each matrix in the sum is skew symmetric, so $U^{-1}\Omega U$ is skew symmetric as well.

Now suppose there is a nonsingular U such that $U^{-1}\Omega U$ is skew symmetric, and that $\Omega = w_1 S_1 + w_2 S_2 + \dots + w_r S_r$, $w_k \in \wedge^2 K$, where the set $\{w_1, w_2, \dots, w_r\}$ is linearly independent. Then

$$U^{-1}\Omega U = \sum_{k=1}^r w_k U^{-1}S_k U \quad (2.4)$$

Since the left hand side of (2.4) is skew symmetric, the right hand side is as well. Consider the (i, j) element of the left hand side of (2.4). This is equal to the negative of the (j, i) element. Hence

$$\sum_{k=1}^r w_k U^{-1}S_k U(i, j) = -\sum_{k=1}^r w_k U^{-1}S_k U(j, i)$$

so, after rearranging, we have

$$\sum_{k=1}^r w_k (U^{-1}S_k U(i, j) + U^{-1}S_k U(j, i)) = 0,$$

and since the w'_k s are linearly independent, this means that $U^{-1}S_k U(i, j) + U^{-1}S_k U(j, i) = 0$ for all i, j . Putting these two things together says that each matrix $U^{-1}S_k U$ is skew symmetric for each k , and that the set S is simultaneously skewable.

□

Let Ω be an $m \times m$ matrix with entries in $\wedge^2 K$, and let $e = (e_1, e_2, \dots, e_n)$ be a frame in K . Then

$$\Omega = \sum_{w_k} w_k S_k, \quad k = 1, 2, \dots, \binom{n}{2} \quad (2.5)$$

where $\{w_k, k = 1, 2, \dots, \binom{n}{2}\} = \{e_i \wedge e_j, 1 \leq i < j \leq n\}$ is the standard basis for $\wedge^2 K$, and the S_k 's have real entries. So we have

Theorem 2.6. Ω as defined above, is skewable if and only if the matrices $\{S_k\}$ are simultaneously skewable.

Next we will give a simple algorithm that can be used to decide whether a set of matrices with real entries are simultaneously skewable.

Lemma 2.7. Let $S = \{S_1, S_2, \dots, S_k\}$ be a set of $m \times m$ matrices with real valued functions as entries. Then S is simultaneously skewable if and only if there exists a positive definite symmetric matrix A , with real valued functions as its entries, such that

$$S_i A + A S_i^T = 0, \quad \forall i. \quad (2.8)$$

If the solution to the system exists, then there is always a symmetric matrix U , with real valued functions as its entries, such that

$$U^2 = A.$$

The matrix U will skew-symmetrize the set S , and its entries are as smooth as the entries of A .

Proof. Suppose that U is a nonsingular real matrix such that for each $1 \leq i \leq m$, $U^{-1} S_i U$ is skew symmetric. Define $A = U U^T$. Clearly A is symmetric and positive definite. We have

$$\begin{aligned} U^{-1}(S_i A + A S_i^T)U &= U^{-1} S_i U U^T U + U^T S_i^T U \\ &= (U^{-1} S_i U) U^T U + (S_i U)^T U \\ &= (U^{-1} S_i U) U^T U + (U U^{-1} S_i U)^T U \\ &= (U^{-1} S_i U) U^T U + (U^{-1} S_i U)^T U^T U \\ &= (U^{-1} S_i U + (U^{-1} S_i U)^T) U^T U \\ &= 0. \end{aligned}$$

Hence $S_i A + A S_i^T = 0$ and the first part is proved.

Conversely, suppose there is a positive definite symmetric matrix A such that $S_i A + A S_i^T = 0$ for each i . Since A is p.d.s., its logarithm $\log A$ is well defined, symmetric and its entries are smooth in the entries of A . Define

$$U = \sqrt{A} = e^{1/2 \log A}.$$

By the properties of the matrix exponential we have

$$U^2 = e^{1/2 \log A} e^{1/2 \log A} = e^{\log A} = A$$

i.e. A has a square root.

We have the following:

$$\begin{aligned} 0 &= S_i A + A S_i^T = U^{-1}(S_i A + A S_i^T)U^{-1} \\ &= U^{-1} S_i U^2 U^{-1} + U^{-1} U^2 S_i^T U^{-1} = U^{-1} S_i U + (U^{-1} S_i U)^T \end{aligned}$$

So $U^{-1} S_i U$ is skew symmetric for each i and the set S is simultaneously skewable. $\square$

Please note that the equations (2.8) define an overdetermined (in general!) system of linear equations with the entries of the matrix A as unknowns.

3 Application to curvature matrices

Let K be 3 dimensional real vector field and let e^1, e^2, e^3 be a frame in K . Consider the "curvature" matrix

$$\Omega = \begin{vmatrix} -e^1 \wedge e^2 - e^3 \wedge e^1 & -2e^1 \wedge e^2 - e^2 \wedge e^3 - e^3 \wedge e^1 & -e^1 \wedge e^2 - 2e^2 \wedge e^3 + e^3 \wedge e^1 \\ e^1 \wedge e^2 + e^3 \wedge e^1 & e^1 \wedge e^2 + e^2 \wedge e^3 + e^3 \wedge e^1 & 2e^2 \wedge e^3 \\ -e^3 \wedge e^1 & -e^2 \wedge e^3 - e^3 \wedge e^1 & -e^2 \wedge e^3 \end{vmatrix}$$

It follows that

$$S_1 = \begin{vmatrix} -1 & -2 & -1 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{vmatrix}, S_2 = \begin{vmatrix} 0 & -1 & -2 \\ 0 & 1 & 2 \\ 0 & -1 & -1 \end{vmatrix}, S_3 = \begin{vmatrix} -1 & -1 & 1 \\ 1 & 1 & 0 \\ -1 & -1 & 0 \end{vmatrix}$$

Using Mathematica, we find one solution to the system $S_i X + X S_i^T = 0$, for $i = 1, 2, 3$, is

$$A = \begin{vmatrix} 3 & -2 & 1 \\ -2 & 2 & -1 \\ 1 & -1 & 1 \end{vmatrix}.$$

A is positive definite symmetric, so we use Mathematica again to find the square root of A , which is:

$$U = \begin{vmatrix} 1.56022 & -0.689101 & 0.301417 \\ -0.689101 & 1.17254 & -0.387684 \\ 0.301417 & -0.387684 & 0.871119 \end{vmatrix}.$$

Note that we have rounded off for readability. This leads to round off errors. To get the results below, use more significant digits.

Then we see (counting for round off) the following:

$$U^{-1} S_1 U = \begin{vmatrix} 0 & -0.87112 & -0.387685 \\ 0.87112 & 0 & -0.301416 \\ 0.387685 & 0.301416 & 0 \end{vmatrix}$$

$$U^{-1} S_2 U = \begin{vmatrix} 0 & -0.086268 & -0.483435 \\ 0.086268 & 0 & 0.871119 \\ 0.483435 & -0.871119 & 0 \end{vmatrix}$$

$$U^{-1} S_3 U = \begin{vmatrix} 0 & -0.483435 & 0.784851 \\ 0.483435 & 0 & 0.387685 \\ -0.784851 & -0.387685 & 0 \end{vmatrix}$$

These are all skew-symmetric, which means the matrix U skewsymmetrizes the set $\{S_1, S_2, S_3\}$, demonstrating the result.

From the point of view of differential geometry, it is important to see whether two matrices that skewsymmetrize the same set of matrices are related. More precisely we will prove the following result

Theorem 3.1. *Let U, V be two matrices with equal determinant that skewsymmetrize $\binom{n}{2}$ linearly independent $n \times n$ matrices $S_1, S_2, \dots, S_{\binom{n}{2}}$, i.e. $U^{-1}S_i U$ and $V^{-1}S_i V$ are skew-symmetric for each i . Then*

$$U = VO,$$

with O orthogonal matrix.

In order to prove Theorem 3.1 we need to establish the following lemma

Lemma 3.2. *Let M_n^s denote the $\binom{n}{2}$ dimensional space of $n \times n$ skew-symmetric matrices and let O be a matrix with determinant equal to one. If for all $X \in M_n^s$*

$$O^{-1}XO \in M_n^s,$$

then O is an orthogonal matrix.

Proof. By the Singular Value Decomposition Theorem we have

$$O = P\Delta L, \tag{3.3}$$

where P and L are orthogonal matrices ($P, L \in O(n)$) and Δ is the diagonal matrix of singular values of O . We will first prove that

$$\Delta^{-1}X\Delta \in M_n^s \quad \forall X \in M_n^s. \tag{3.4}$$

Assume the contrary, that there exist a skew symmetric matrix X such that

$$\Delta^{-1}X\Delta \notin M_n^s. \tag{3.5}$$

It follows that

$$\Delta^{-1}P^T PXP^T P\Delta \notin M_n^s, \tag{3.6}$$

and consequently

$$L^T \Delta^{-1}P^T PXP^T P\Delta L \notin M_n^s, \tag{3.7}$$

which in turn means

$$O^{-1}PXP^T O \notin M_n^s \tag{3.8}$$

for the skew symmetric matrix PXP^T . Thus we contradict the hypothesis. It follows that equation (3.4) is true. Applying equation (3.4) to the special frame of $\binom{n}{2}$ linearly independent skew symmetric matrices that have only two non zero entries we conclude that

$$\Delta = I.$$

The conclusion of the lemma follows. □

Now we will prove Theorem 3.1.

Proof. Let U, V be two matrices with equal determinant that skew-symmetrize $\binom{n}{2}$ linearly independent matrices $S_1, S_2, \dots, S_{\binom{n}{2}}$. Let us denote by S the $\binom{n}{2}$ dimensional span of $S_1, S_2, \dots, S_{\binom{n}{2}}$ and let us note that both U and V induce the following linear isomorphisms

$$\begin{aligned} \overline{U} : S &\rightarrow M_n^s \\ \overline{U}(X) &= U^{-1}XU \end{aligned} \tag{3.9}$$

and

$$\begin{aligned}\bar{V} : S &\rightarrow M_n^s \\ \bar{V}(X) &= V^{-1}XV\end{aligned}\tag{3.10}$$

These two mappings are clearly linear and injective and by reasons of dimension it follows that both are indeed linear isomorphisms. It follows that

$$\bar{U} \circ \bar{V}^{-1} : M_n^s \rightarrow M_n^s$$

is a linear isomorphism defined by the equation

$$\bar{U} \circ \bar{V}^{-1}(X) = U^{-1}VXV^{-1}U\tag{3.11}$$

Since

$$\det U^{-1}V = 1$$

and by applying Lemma 3.2 we conclude that $\exists O \in \mathcal{O}(n)$ such that

$$U = VO.$$

□

As a consequence of Theorem 3.1 we will be able to construct an algorithm that will allow us to verify whether a connection with full rank curvature matrix is locally metric. First we will clarify what we mean by full rank curvature.

Definition 3.12. Let D be connection in the vector bundle E . Let

$$\sigma = (\sigma_1, \sigma_2, \dots, \sigma_m)$$

be a local frame in the contractible open set $N \subset M$. Let Ω be the curvature matrix with respect to σ . Consider

$$(\theta^i \wedge \theta^j)_{1 \leq i < j \leq n}$$

be a local frame of two forms on N . Let S_{ij} be the matrices with entries smooth, real valued functions on N defined by the equation

$$\Omega = \sum_{1 \leq i < j \leq n} (\theta^i \wedge \theta^j) S_{ij}\tag{3.13}$$

The number r of linearly independent matrices $(S_{ij})_{1 \leq i < j \leq n}$ will be called the rank of Ω .

If

$$r = \binom{n}{2}$$

then Ω is said to have full rank.

It is trivial to verify that the rank of the curvature is a pointwise, scalar invariant of the connection.

In order to justify the algorithm, we need the following corollary of Theorem 3.1:

Corollary 3.14. Let D be a connection in a vector bundle. Assume that the dimension of its fibers is equal to the dimension of the base manifold (i.e. $m = n$). Assume D is metric on a contractible neighborhood $N \subset M$, the curvature matrix Ω is skew symmetric with respect to a local frame

$$\sigma = (\sigma_1, \sigma_2, \dots, \sigma_n),$$

and it has full rank on N . Then the connection matrix ω with respect to σ is skew-symmetric as well.

Proof. Let

$$(\theta^i \wedge \theta^j)_{1 \leq i < j \leq n}$$

be a local frame of two forms on N , and S_{ij} be the family of matrices defined by equation (3.13). Because Ω is skew and of full rank, it follows that S_{ij} 's are linearly independent and also skew and therefore skew-symmetrized by the identity matrix I . Since D is assumed locally metric, then there exist a frame $\tau = (\tau_1, \tau_2, \dots, \tau_n)$ such that the connection matrix ψ with respect to this frame is skew and consequently the curvature matrix Ψ . Let B be the matrix defined by

$$\tau = \sigma B.$$

We have

$$\Psi = B^{-1} \Omega B = \sum_{1 \leq i < j \leq n} (\theta^i \wedge \theta^j) B^{-1} S_{ij} B$$

and according to Lemma 2.3 it follows that S_{ij} 's are skew-symmetrized by the matrix B . According to Theorem 3.1 it follows that

$$B = IO = O$$

with O an orthogonal matrix. Hence B is orthogonal.

Let ω be the connection matrix with respect to σ . It follows that

$$\psi = B^T dB + B^T \omega B.$$

But since B is orthonormal, from

$$B^T B = I$$

(by differentiating) it follows that

$$B^T dB \text{ is skew.}$$

Hence

$$B^T \omega B \text{ is skew}$$

and therefore ω is skew. □

Now consider a connection D in the vector bundle E . Assume that the dimension of its fibers is equal to the dimension of the base manifold (i.e. $m = n$). We assume that the curvature has full rank on a contractible open set $N \subset M$. We would like to verify if the connection is metric on N .

(a) Take a local frame $\sigma' = (\sigma'_1, \sigma'_2, \dots, \sigma'_m)$ and calculate the curvature matrix Ω' with respect to this frame.

(b) Determine the matrices S'_{ij} defined by Ω' as in equation (3.13).

(c) Form the system

$$X S'_{ij} + (S'_{ij})^T X = 0.$$

(d) If the linear system from the previous step does not have a symmetric, positive definite solution $X = A$, then D is not metric.

(e) If the solution A exists then take its square root $\sqrt{A} = U^{-1}$.

(f) Form the frame σ defined by

$$\sigma = \sigma' U^{-1}.$$

The curvature matrix Ω with respect to σ is skew-symmetric. Calculate the connection matrix ω with respect to σ and check if it is skew. If it is not, then the connection is not metric on N according to Corollary 3.14.

- (g) If the connection matrix ω with respect to σ is skew, then the metric that makes σ orthonormal is compatible with the connection.
-

REFERENCES

- [1] Theodore Frankel, "The geometry of physics. An Introduction", Cambridge University Press (2011)
- [2] Mihail Cocos, "A note on locally metric connections", arXiv:1606.05813 [math.DG]
- [3] Alena Vanžurová, Petra Žáčková, "Metrizability of connections on two-manifolds", Acta Universitatis Palackianae Olomucensis. Facultas Rerum Naturalium. Mathematica , Vol. 48 (2009), No. 1, 157–170
- [4] Vanžurová, A., Žáčková, P., Metrization of linear connections, Aplimat, J. of Applied Math. (Bratislava) 2, 1 (2009), 151163.
- [5] Thompson, G.: "Local and global existence of metrics in two-dimensional affine manifolds", Chinese J. Phys. 19, 6 (1991), 529532.
- [6] Cheng, K. S, Ni, W. T.: "Necessary and sufficient conditions for the existence of metrics in two-dimensional affine manifolds", Chinese J. Phys. 16 (1978), 228232.
- [7] Oldrich Kowalski, Martin Belger, "Metrics with the Prescribed Curvature Tensor and all Its Covariant Derivatives at One Point", Mathematische Nachrichten Volume 168, Issue 1, pages 209225, 1994