Comparative Effectiveness of General Anesthesia in Doves Using a Combination of Ketamine and Diazepam

Syed Wasim Abbas^{1*}, Muhammad Naveed Ali¹, Ghulam Abbas², Hafiz Muhammad Safwan¹,
Muhammad Sajid³, Memoona Mehmood⁴

¹Department of Clinical Medicine and Surgery, University of Veterinary and Animal Sciences, Pakistan

²Department of Animal Production, Riphah College of Veterinary Sciences, Riphah International University, Pakistan

³Department of Anatomy, Riphah College of Veterinary Sciences, Riphah International University, Pakistan

⁴Faculty of Veterinary Science, University of Agriculture Faisalabad, Pakistan

Copyright©2018 by authors, all rights reserved. Authors agree that this article remains permanently open access under the terms of the Creative Commons Attribution License 4.0 International License

Abstract The present research was conducted to check the clinical effects of ketamine, diazepam and a ketamine and a combination of diazepam in the general anaesthesia of doves. 32 doves of both sexes with body weights ranging from 280 gm to 300 gm were divided randomly to 4 groups having 8 birds each. Group A received a 0.5 mL mixture of diazepam (0.2 mg/kg) and normal saline, group B a 0.5 mL mixture of ketamine 5% (30 mg/kg) and normal saline, group C a 0.5 mL mixture of ketamine 5% (10 mg/kg), diazepam (0.2 mg/kg) and normal saline, while group D (control) received 0.5 mL of normal saline only. Each mixture was administered intramuscularly. Under standard operating room conditions, general anaesthesia was not observed in group D (normal saline alone). In group A, sedation and muscle relaxation without complete loss of consciousness was observed. Induction time of anaesthesia in group C (doves treated with ketamine and diazepam combination) was significantly quicker than group B ($p < 0.05$) that were treated with ketamine alone. Duration of anaesthesia in group C was significantly longer than group B ($p < 0.05$). Recovery took longer in group C in comparison with group K, but the difference was not statistically significant ($p > 0.05$). The birds in group C were calm and sedated, with good muscle relaxation, whilst in group B the birds were excited and showed a drop in body temperature. According to the results of this study, the combination of low dose ketamine hydrochloride (HCL) and diazepam overcame the adverse effects of ketamine alone. This combination produced a more rapid induction of anaesthesia, as well as an increase in anaesthesia duration, with good muscle relaxation and a smooth and slow recovery. Use of a combination of ketamine HCL given at 10 mg/kg and diazepam given at 0.2 mg/kg for anaesthesia in doves is therefore recommended.

Keyword Effects, Ketamine, Diazepam, Anaesthesia, Doves

1. Introduction

Avian anesthetic and surgical techniques have advanced vastly in the last decade. The selection of anesthesia and route of administration is often as essential as the surgical procedure itself [1]. General anaesthesia in different avian species may be produced by administration of either inhalation or inject-able agents [2;3]. Injectable anaesthetics and sedatives can be administered intramuscularly, intravenously or intraosseously. [4] Inhalation of anaesthesia is preferred for birds but needs costly equipment. The use of an injectable agent in comparison with an inhalation agent has the advantage of increased speed of anaesthesia induction, the need for minimal equipment and a minimum cost [2;3;5].

Several injectable drugs are used in birds. These contain barbiturates, chloral hydrate, phenothiazine derivatives, alpha-2 adrenergic agonists, ketamine and propofol [6; 7;8]. Pigeons are very sensitive birds and any mismanagement in a crisis can induce direct concussion and death. Doves are often referred to a hospital in a critical condition and these birds need safe anaesthesia and painless surgery. In these circumstances, careful selection of the safest option of anaesthetic agent is very essential [9; 10]. Ketamine is a dissociative anaesthetic agent which can be employed for the induction of general anaesthesia in many species by either the intravenous, intramuscular or intraosseous route [1; 5; 6]. However, ketamine is seldom used as the only sedative agent in birds. According to Athar et al. [2], ketamine has a broad safety margin in birds and as much as 10 times the therapeutic dose is normally required before symptoms of toxicity appear. For respiratory depression that may result from ketamine, supportive ventilation and administration of doxapram can be applied.

When ketamine is used as the solitary anaesthetic agent, it induces poor muscle relaxation, muscle tremors, myotonic contractions, opisthotonos, persistent pain reflexes and stormy recoveries [11; 12; 13]. Therefore, it is most often combined with other agents (alpha-2 adrenergic agonists, diazepam or azaperone), depending on the species involved [10; 14; 15]. A combination of ketamine and benzodiazepines (diazepam and midazolam) or ketamine and alpha-2 adrenergic agonists are commonly used for general anaesthesia in pigeons [9; 10; 15; 16]. Diazepam has potent muscle relaxant and anticonvulsant properties and has been used in a vast range of wild and domestic animals and birds [15; 16 17]. When used together, diazepam and ketamine have a synergistic effect, resulting in a smooth recovery and better muscle relaxation [11; 18]. Their efficacy is enhanced whilst minimising their unwanted adverse effects.

The objective of this research was to compare the effects (time to onset of anaesthesia, duration of anaesthesia, duration of recovery, response to external stimuli, muscular relaxation, palpebral and pedal reflexes, and cloacal temperature) of diazepam and ketamine in doves when administered intramuscularly alone or combined.

2. Materials and Methods

Birds

Thirty-two healthy adult doves of both sexes (23 males and 9 females) with body weights ranging from 280 g to 300 g were used in the study. All the doves were from the same flock and their ages were between 1 and 2 years. All the birds were acclimatised in a calm room for 2 weeks. They were fed a wheat-based diet similar to their previous diet and had free availability of water and food, except for a period of 1 h prior to drug administration. This decreased the occurrences of vomiting. Before the start of the study, the birds were found to be in good health, based on physical examinations.

Trial Procedure

The research was conducted at the Department of Clinical Medicine and Surgery, University of Veterinary and Animal Sciences, Pakistan. For this, doves were placed randomly to four equal groups of eight birds each. The groups were differentiated as follows: in group A, diazepam was administered at a dose of 0.2 mg/kg, in group B, ketamine 5% was administered at a dose of 30 mg/kg, in group C, a ketamine (5%) and diazepam combination was administered at a dose of 10 mg/kg and 0.2 mg/kg, respectively, whilst in group D (control), a 0.5 mL dose of normal saline was given per bird.

All medications were administered intramuscularly into the deep pectoral muscle using an insulin syringe. The medications in groups A, B and C were diluted with normal saline to make a final volume of 0.5 ml.

Post-treatment Monitoring

After drug administration, each bird was placed in sternal recumbency in separate cages, so that each bird could be observed individually and external stimulation was kept to a minimum.

The clinical parameters for each dove were as follows:

- The severity of opisthotonos was scored on a scale of 0–4: lack of opisthotonos was scored as 0 and those with severest opisthotonos were scored as 4.
- Onset of anaesthesia: Time interval (in minutes) from administration through the stages of anaesthesia to loss of consciousness.
- Duration of anaesthesia: Time interval (in minutes) from loss of consciousness to reappearance of sensation.
- Eyelids: scored as - for closed, + for half-opened and ++ for opened.
- Duration of recovery: Time interval (in minutes) from the return of reflexes to complete consciousness.
- The severity of excitability, such as wing flapping and ataxia was scored from 0 to 1. No excitability was scored as 0 and the birds showing most excitability were scored as 1.
- Muscle relaxation was tested in the muscles of the neck, wings and legs. The ease with which the wings of the birds could be pulled, their hind limbs could be flexed and the neck could be extended was recorded as the degree of muscle relaxation. It was graded on a scale of 0.0–3.0, where 0.0 represented weak relaxation. Birds given this score showed closed wings and stiff limbs. A score of 1.5 represented moderate relaxation where there was mild resistance to extension of the neck, opening of the wings and flexing of the limbs, whilst a score of 3.0 represented excellent relaxation where the birds showed a flaccid neck, no resistance to opening of the wings and flexing of limbs.
- Pedal reflex was evaluated by using a towel clamp forceps. The presence and strength of the reflex was scored based on a scale of 0–4, with 0 being a weak reflex and 4 being a strong reflex.
- Palpebral reflexes were tested by touching the eyelids with a sterile cotton tip swab. Lack or presence of the reflexes were scored as - or +, respectively.

Cloacal temperature was measured before anaesthesia and again 10 min after induction of anaesthesia. A digital thermometer was used.

Statistical Analysis

Opisthotonos, onset of anaesthesia, duration of anaesthesia and duration of recovery were compared using an independent *t*-test. The Kruskal-Wallis test was used to compare the mean scores amongst the different groups. Cloacal temperature was analysed by one-way ANOVA

and Duncan's multiple-range test. All statements of significance were based on the 0.05 level of probability.

Ethical Considerations

The Institutional Animal Care and Use Committee approved the protocol for this project, under the following project number 671.

3. Results

No anaesthesia or opisthotonos was observed in groups D and A. Diazepam alone showed better muscle relaxation than ketamine alone, but the muscle relaxation was not as good as with the ketamine and diazepam combination. Opisthotonos in the ketamine and diazepam combination was more severe than with ketamine alone, but the difference was not statistically significant ($p > 0.05$) (Table 1). The onset of anaesthesia after injection was significantly quicker with the ketamine and diazepam combination ($1.50 \text{ min} \pm 0.23 \text{ min}$) than with ketamine

alone ($4.50 \text{ min} \pm 0.41 \text{ min}$) ($p < 0.05$) (Table 1). The duration of anaesthesia in the C group ($14.10 \text{ min} \pm 1.48 \text{ min}$) was significantly higher than group B, where ketamine was used alone ($8.13 \text{ min} \pm 1.41 \text{ min}$) ($p < 0.05$) (Table 1). Recovery from anaesthesia took longer with the ketamine and diazepam combination ($23.12 \text{ min} \pm 2.85 \text{ min}$) compared with ketamine alone ($19.25 \text{ min} \pm 2.69 \text{ min}$), but the difference was not statistically significant ($p > 0.05$) (Table 1). The excitability during the recovery was significantly less in group C as compared with group B ($p < 0.05$). The pigeons that were treated with diazepam alone were calm and sedated during recovery.

Muscle relaxation following the administration of the ketamine and diazepam combination was twice as pronounced than with diazepam alone ($p < 0.05$). Muscle relaxation was not observed in groups B and D (Figure 1). The pigeons' pedal reflexes were significantly weaker in groups B and C when compared with groups D and A ($p < 0.05$). This reflex was markedly weaker with the ketamine and diazepam combination than with ketamine alone (Figure 2).

Table 1. Rate of opisthotonos and time taken in minutes for onset of anaesthesia, duration of anaesthesia and recovery period in doves treated with ketamine only and a ketamine diazepam combination.

Groups	Ophisthotonos	Onset of anaesthesia	Duration of anaesthesia	Recovery period	Excitement
B	3.75 ± 0.31	4.37 ± 0.21^a	8.55 ± 1.48^a	19.67 ± 2.69	0.85 ± 0.05^a
C	3.85 ± 0.39	1.89 ± 0.23^b	14.52 ± 1.46^b	23.77 ± 2.85	0.15 ± 0.04^b

B, doves treated with ketamine; C, doves treated with ketamine and diazepam combination.

^{a,b}, mean presented in column with different superscripts differing significantly ($P < 0.05$).

NS, normal saline; K, ketamine; D, Diazepam; KD, ketamine diazepam

Figure 1. The comparison of muscular relaxation across the groups of doves treated with normal saline, ketamine, diazepam, and a ketamine and diazepam combination, where a score of 0.0 represents weak relaxation, 1.5 represents moderate relaxation and 3.0 represents excellent relaxation.

NS, normal saline; K, ketamine; D, Diazepam; KD, ketamine diazepam

Figure 2. The comparison of pedal reflex across the groups of doves treated with normal saline, ketamine, diazepam, and a ketamine and diazepam combination, where a score of 0.0 represents a weak reflex and 4.0 represents a strong reflex.

There were significant differences in eyelid status between group C and groups D, A and B ($p < 0.05$). The eyelids of pigeons treated with the ketamine and diazepam combination were completely closed. With ketamine alone, the pigeons' eyes were completely open during anesthesia (Table 2). There were no significant differences in the palpebral reflexes in the different groups and these were present in all the groups (Table 2). The cloacal temperature was $41.30^{\circ}\text{C} \pm 0.72^{\circ}\text{C}$ before medication. When measured 10 min after drug administration in groups B and A, the cloacal temperature was recorded to have decreased significantly ($p < 0.05$). However, in pigeons from group C, the cloacal temperature remained within the initial range, measuring at $40.93^{\circ}\text{C} \pm 0.86^{\circ}\text{C}$ (Table 2).

Table 2. The status of eyelids, response of palpebral reflex and cloacal temperature in doves of different groups

Groups	Eyelids	Palpebral Reflex	Cloacal Temperature
A	++	+	38.52 ± 0.04^b
B	++	+	39.17 ± 0.59^b
C	-	+	40.93 ± 0.86^a
D	++	+	41.30 ± 0.72^a

A, doves treated with diazepam; B, doves treated with ketamine; C, doves treated with ketamine and diazepam combination; D, doves treated with normal saline;

++, eyelids open; -, eyelids closed; +, palpebral reflex present.

^{a,b}, mean presented in column with different superscripts differing significantly ($P < 0.05$).

4. Discussion

The dose and anaesthetic response of ketamine varies across different bird species.[13] The recommended dosage of ketamine is approximately 10 mg/kg – 60 mg/kg, given intramuscularly or intravenously. Larger birds (greater than 1000 g) require a lower dose per kilogram (10 mg/kg). The recommended intramuscular and/or intravenous dosage of diazepam in birds is 0.2 mg/kg – 1 mg/kg.[11] In this study, the dosage of ketamine was about half the normal dose used in pigeons.[9,10,15] There are very few articles in the current literature on the ideal intramuscular dosage of the diazepam and ketamine combination in pigeons. For this reason a low dosage of ketamine and diazepam was used.

A ketamine and diazepam combination produced a fast and smooth induction of anaesthesia, whilst a dosage of ketamine alone produced a slow and smooth anaesthesia. The duration of anaesthesia with a ketamine and diazepam combination ($14.10 \text{ min} \pm 1.48 \text{ min}$) obtained in this study was longer than the $8.13 \text{ min} \pm 1.41 \text{ min}$ obtained with ketamine alone. Diazepam is distributed widely throughout the body after administration and, being fat soluble, it can be deposited into muscle and fat tissue.¹⁹ The diazepam that was absorbed by the fat and muscle tissue is then released slowly into the system, resulting in a longer duration of anaesthesia than with ketamine alone.

Paul-Murphy et al. [20] reported that the average surgical time needed for most perching bird species was 15 min. Therefore, the increase in anaesthesia duration through this combination provides enough time for most surgical procedures. Ketamine elicited hyperthermia, tachycardia and hypoventilation during anaesthesia. Similar to the observations of Virtanen [21] and Sandmeier [22], detomidine – ketamine cocktail synergistically produced a deep anaesthesia accompanied by slight hypothermia, hypoventilation and bradycardia. Diazepam is highly lipid-soluble, and is widely distributed throughout the body after administration and it can redistribute into muscle and adipose tissue [19]. In avian species 2 techniques are recommended for sedation and anaesthesia: inhalational and parenteral. The inhalational technique typically involves the use of isoflurane or sevoflurane, while parenteral methods utilise intravenous, intramuscular, or subcutaneous administration of injectable drugs (xylazine, medetomidine, detomidine, ketamine, chloral hydrate, barbiturates, butorphanol, and phenothiazines). Freed and Baker [23] reported that xylazine, detomidine, and medetomidine are usually combined with ketamine in birds and administered intramuscularly or intravenously. The alpha-2-adrenergic agonists provide muscle relaxation, analgesia, and sedation and smooth induction and recovery. They suppress pain arising superficially and also visceral pain.

The xylazine-ketamine cocktail gave a very fast and smooth induction of anaesthesia associated with deep analgesia, good muscle relaxation, and absence of all reflexes. These findings are in agreement with the findings published by Krajka and Juranova [7]. Shakoor et al. [25] also noted similar hypothermia, hypoventilation, and bradycardia throughout anaesthesia until complete recovery. Recovery was smooth in all our birds with some efforts to attain their normal sitting or standing posture. The same observations are reported by Mohammad et al. [24], who studied the effect of detomidine in combination with ketamine in chickens.

In the present study, the recovery was smooth but slow in the pigeons anaesthetised with the ketamine and diazepam combination. This observation is similar to those reported by Lumeij and Deenik [15] and Varner et al [18]. In the ketamine-treated pigeons, recovery was stormy. These birds showed severe convulsions and wing flapping. This was also found by Atalan et al. [14], who reported that the stormy recovery is caused by the dissociative characteristics of ketamine anaesthesia. With the ketamine and diazepam combination, the reduction of side effects during the recovery could be the result of the low dose of ketamine used in this group, as compared with ketamine alone (used at a higher dosage), as well as the simultaneous administration of sedatives such as diazepam.

In this study, muscle relaxation was significantly more pronounced in the ketamine and diazepam combination than with diazepam alone ($p < 0.05$). Improvement of

muscular relaxation in this combination was associated with the muscle relaxant properties of diazepam [15; 18; 19]. Poor muscle relaxation was found when ketamine was used alone. [13; 14]. The presence of palpebral reflex in the ketamine-treated pigeons was reported previously by many studies [9; 10] and was also observed in our study. In the present study, the cloacal temperature 10 min after anaesthetic administration dropped to $39.17^{\circ}\text{C} \pm 0.59^{\circ}\text{C}$ in the ketamine group. Yet, it remained at $40.93^{\circ}\text{C} \pm 0.86^{\circ}\text{C}$ in pigeons treated with the ketamine and diazepam combination.

The present study showed that intramuscular injection of the diazepam and normal saline mixture could not induce anaesthesia, but sedation and muscle relaxation were observed. Normal saline alone produced no sedation, muscle relaxation or anaesthesia. Solution volume alone therefore plays no role. This finding correlates with previous investigations [15; 18].

5. Conclusions

The combination of ketamine HCL given at 10 mg/kg and diazepam given at 0.2 mg/kg provides more ideal conditions for the veterinary surgeon. This is, to a large extent, thanks to the sedation and the prolongation of the anaesthetic effect produced by diazepam. The ketamine and diazepam combination produced more a rapid induction of anaesthesia, an increase in the duration of anaesthesia, as well as smooth recovery and more muscle relaxation without any adverse effects. It can therefore be considered an important tool for the light anaesthesia of pigeons.

REFERENCES

- [1] Forbes, N. A. 1998. Avian anesthesia. *Vet Q.* 20:265–266.
- [2] Athar M, Shakoor A, Muhammad G, Sarwar MN, N. I. Chaudhry. 1996. Clinical perspectives of intravenous ketamine anesthesia in peafowl (*Pavo cristatus*). *Acta Vet Hung.* 44:357–361.
- [3] Christense, J., Fosse R. T, Halvorsen O. J, Moril I. 1987. Comparison of various anesthetic regimens in the domestic fowl. *Am J Vet Res.*, 48:1649–1657.
- [4] Harrison, GJ. Anesthesiology. In: Harrison J, Harrison LR, editors. *Clinical avian medicine and surgery*. 7th edn. Philadelphia: WB Saunders, 1986; p. 520–545.
- [5] Samour JH, Jones DM, Knight JA, Howlett JC. Comparative studies of the use of some injectable anesthetic agents in birds. *Vet Rec.* 1984; 115:6–11.
- [6] Hall LW, Clarke KW, Trim CM. Principles of sedation, analgesia and premedication. In: Hall LW, Clarke KW, Trim CM, editors. *Veterinary anesthesia*. 10th edn. Oxford: Elsevier Ltd, 2001; p. 292–316.
- [7] Krajka A, Juranova R. Anesthesia in poultry. *Vet Med.* 1994; 39: 23–27.
- [8] Machin KL, Caulkett NA. Cardiopulmonary effects of propofol and a medetomidine-midazolam-ketamine combination in Mallard ducks. *Am J Vet Res.* 1998; 59: 598–602.
- [9] Durrani UF, Arif khan M, Saleem ahmad S. Comparative efficacy (sedative and anesthetic) of detomidine, ketamine and detomidine, ketamine cocktail in pigeons (*Columba livia*). *Pakistan Vet J.* 2008; 28: 115–118.
- [10] Durrani UF, Ashraf M, Khan MA. A comparison of the clinical effects associated with xylazine, ketamine, and a xylazine, ketamine cocktail in pigeons (*Columba livia*). *Turk J Vet Anim Sci.* 2009; 33: 413–417.
- [11] Patrick R. Anesthesia, analgesia and monitoring. Paper presented at: The North American Veterinary Conference; 2005 Jan 08–12; Orlando, USA.
- [12] Valverde A, Bienzle D, Smith DA, Dyson DH, Valliant AE. Intraosseous cannulation and drug administration for induction of anesthesia in chickens. *Vet Surg.* 1993; 22: 240–244.
- [13] Wright M. Pharmacologic effects of ketamine and its use in veterinary medicine. *J Am Vet Med Assoc.* 1982; 180:1462–1470.
- [14] Atalan G, Uzun M, Demirkan S, Yildiz S, Cenesiz M. Effect of medetomidine- butorphanol-ketamine anesthesia and atipamezole on heart and respiratory rate and cloacal temperature of domestic pigeons. *J Vet Med A Physiol Pathol Clin Med.* 2002; 49: 281–285.
- [15] Lumeij JT, Deenik JW. Medetomidine, ketamine and diazepam-ketamine anesthesia in racing pigeons (*Columba livia domestica*): A comparative study. *J Avian Med Surg.* 2003; 17: 191–196.
- [16] Zamani Moghadam A, Bigham Sadegh A, Sharifi S, Habibian S. Comparison of intranasal administration of diazepam, midazolam and xylazine in pigeons: Clinical evaluation. *Iran J Vet Sci Technol.* 2009; 1: 19–26.
- [17] Lees P. Sedatives. In: Lees P, editor. *Veterinary applied pharmacology and therapeutics*. 5th edn. Oxford: Baillière Tindall, 1991; p. 337–339.
- [18] Varner J, Clifton KR, Poulos S, Broderson JR, Wyatt RD. Lack of efficacy of injectable ketamine with xylazine or diazepam for anesthesia in chickens. *Lab Anim.* 2004; 33:36–39.
- [19] Bateson AN. Basic pharmacologic mechanisms involved in benzodiazepine tolerance and withdrawal. *Curr Pharm Des.* 2002;8:5–21.
- [20] Paul-Murphy JR, Brunson DB, Miletic VA. Technique for evaluating analgesia in conscious perching birds. *Am J Vet Res.* 1999;60:1213–1217.
- [21] Virtanen, R., 1986. Pharmacology of detomidine and other alpha-2-adrenoceptor agonists in the brain. *Acta Vet. Scand. Suppl.*, 82: 35–46.
- [22] Sandmeier, P., 2000. Evaluation of medetomidine for short term immobilization of domestic pigeons (*Columba livia*) and Amazon parrot (*Amazona species*). *J. Avian Med. Surg.*, 14(1): 8–14.

- [23] Freed, D., Baker, B.: Antagonism of xylazine hydrochloride sedation in raptors by yohimbine hydrochloride. *J. Wildl. Dis.*, 1989; 25: 136-138.
- [24] Mohammad, F.K., al-Badrany, M.S., al-Hassan, A.M.: Detomidine-ketamine anaesthesia in chickens. *Vet. Rec.*, 1993; 133: 192.
- [25] Shakoor, A., Athar, M., Muhammad, G., Chaudhry, N.I.: Anaesthesia in avian species. I: Study on the efficacy of two dose rates of ketamine in peafowls. *Pak. Vet. J.*, 1996; 16: 140- 143.