

Matter from Toric Geometry and its Search at the LHC

T.V. Obikhod

Institute for Nuclear Research, NAS of Ukraine 03680 Kiev, Ukraine

*Corresponding Author: obikhod@kinr.kiev.ua

Copyright ©2013 Horizon Research Publishing All rights reserved.

Abstract Toric geometry is applied for construction the enhanced gauge groups in F-theory compactified on elliptic Calabi-Yau fourfolds. The Hodge numbers calculated from the polyhedra for the chain $H = SU(1), \dots, SU(5), SO(10), E_6, E_7$ determine the number of tensor multiplets, vector multiplets and hypermultiplets of solitonic states that appear from singularities of elliptic fibration. Due to duality between the compactification of E_8 times E_8 heterotic string and the type IIA string compactification on a Calabi-Yau manifold there is a natural sequence of E-group embeddings which gives the matter content of Minimal Supersymmetric Standard Model and the possibility of searching for supersymmetry at the LHC.

Keywords Toric geometry, Calabi-Yau manifold, Singularities of elliptic fibration, Matter content

1 Introduction

At high energy, the three gauge interactions which define the electromagnetic, weak, and strong interactions, are merged into one single interaction characterized by one larger gauge symmetry Grand Unified Theory, (GUT) [1] and thus one unified coupling constant. This gauge coupling unification works quite well in the Minimal Supersymmetric Standard Model (MSSM) motivating not only grand unification (or some superstring theories with similar properties) but also that supersymmetry emerges. Supersymmetry refers to possible relations between the spectrum and interactions of fermions (half-integer spin particles) and bosons (integer spin particles). It can be viewed as a space-time extension of the Poincare (Lorentz plus translational invariance) group, involving new anti-commuting dimensions. Under reasonable assumptions it is the unique extension of the usual Poincare and internal symmetries of field theory. Supersymmetry provides a possible route to unify gravity with the other interactions through superstring theory [2]. For a consistent formulation superstring theories require additional dimensions and there are actually a large number of string theories, many of which include underlying grand unification symmetries (GUT). One of the branches of string theory is F-theory [3] which allows string theorists to construct new realistic vacua - in the form of F-theory compactified on elliptically fibered Calabi-Yau fourfolds.

A coherent description of the fundamental interactions at high energies is the first motivation for the appearance of extra dimensions [4]. In the case of superstring theory, consistency requires spacetime to have 10 (3+1+6) dimensions. If the extra dimensions are compactified, then the extra six dimensions must be in the form of a Calabi-Yau manifold. Compactification on Calabi-Yau n-folds are important because they leave some of the original supersymmetry unbroken. It is natural to consider F-theory compactifications on a Calabi-Yau threefold which admits an elliptic fibration [3]. We will study elliptic Calabi-Yau manifolds which in addition admit a K3 fibration; in other words we consider the case where the K3 fiber itself is elliptically fibered. This would be useful for dualities with heterotic strings. Then the proposed Calabi-Yau must have a singularity. Our purpose is to describe a general method for constructing of Calabi-Yau hypersurfaces in toric varieties. From the toric description of the Calabi-Yau manifold used for compactification can be read off the Newton polyhedron that provides nonabelian gauge groups occurring in type IIA and F-theory. There is a natural sequence of group embeddings which gives the Minimal Supersymmetric Standard Model gauge group and matter structure that can be detected at the LHC [5].

2 Reflexive Polyhedra as Geometric Realization of Calabi-Yau Fourfolds

The elegant algorithm for obtaining the enhanced gauge groups in F-theory has been proposed by Candelas et al. [6, 7]. We consider F-theory of the form $R^{3,1} \times X$ [8, 9], where X is a Calabi-Yau fourfold. We consider superstring compactification on the Calabi-Yau fourfold X represented as the elliptic fibration over a complex

threefold - the Hirzebruch surface F_n , $\mathcal{E} \rightarrow X$
 \downarrow We also take X to be a local elliptic $K3$ -fibration over
 F_n
 $Y \rightarrow X$
 \downarrow where we model the local elliptically-fibered $K3$ -surface Y on a hypersurface in
 S
 \downarrow
 S

\mathbb{C}^3 with an isolated ADE singularity. A great many Calabi-Yau manifolds are known in terms of toric data [10] and there is the correspondence, pointed out by Batyrev, between Calabi-Yau manifolds and reflexive polyhedra Δ [11]. To a Calabi-Yau manifold (of any dimension) defined as a hypersurface in a weighted projective space one can associate its Newton polyhedron, which we denote by Δ . The Newton polyhedron is often reflexive and when it is we may define the dual or polar polyhedron which we denote by ∇ . Polyhedra Δ and ∇ are reflexive, when fourfold X is simultaneously $K3$ -fibration and can also be viewed as an elliptic fibration over the Hirzebruch surface F_{2k} . By means of a computer program we have computed the dual polyhedra for the fourfolds $X_{18k+18}(1, 1, 1, 3k, 6k+6, 9k+9)(k=1, \dots, 6)$.

The following observations summarize the structure of the polyhedron ∇ :

1. Omitting the first two points and the last point of ∇ leaves us with the dual polyhedron ${}^3\nabla$ of the $K3$ surface $X_{12}(1, 1, 4, 6)$;
2. Omitting the first three points and the last two points of ∇ leaves us with the dual polyhedron ${}^2\nabla$ of the torus $X_6(1, 2, 3)$;
3. The polyhedron ${}^3\nabla$ is divided into a ‘top’ and a ‘bottom’ by the polyhedron ${}^2\nabla$ and we may write

$${}^3\nabla = \nabla_{bot}^H \cup \nabla_{top}^{k=1},$$

where $\nabla_{top}^{k=1}$ depends only on $k=1$ while ∇_{bot}^H depends only on the enhanced gauge group H .

Let us introduce special points $pt_1^{(j)} = (0, 0, -j, 2, 3)$,

$$pt_2^{(j)} = (0, 0, -j, 1, 2),$$

$$pt_3^{(j)} = (0, 0, -j, 1, 1),$$

$$pt_4^{(j)} = (0, 0, -j, 0, 1),$$

$$pt_5^{(j)} = (0, 0, -j, 0, 0),$$

$$pt_6^{(j)} = (0, 0, -j, -1, 0),$$

$$pt_7^{(j)} = (0, 0, -j, 0, -1),$$

where j is a positive integer.

We consider now the possibility of adding to ${}^2\nabla$ combinations of the points $pt_r^{(j)}$ in all possible ways such that the bottom corresponds to a reflexive polyhedron. Tables 1 and 2 show the allowed bottoms leading to enhanced gauge groups H . Note that in each case the points of ${}^2\nabla$ are understood and $pt_r^{(j)}$ implies the presence of $pt_r^{(j-1)}, \dots, pt_r^{(1)}$.

Applying the analogous consideration to tops we obtain the following gauge content for fourfolds (1) :

$$\begin{array}{ll} H \times SU(1) & \text{for } k=1, \\ H \times SO(8) & \text{for } k=2, \\ H \times E_6 & \text{for } k=3, \\ H \times E_7 & \text{for } k=4, \\ H \times E_8 & \text{for } k=5, \\ H \times E_8 & \text{for } k=6. \end{array}$$

The relations between the bottoms and the enhanced gauge groups are presented in Table 1.

3 Nesting of polyhedra

We consider now the possibility of adding combinations of the points in all possible ways such that the bottom corresponds to a reflexive polyhedron. All the groups do appear in Table 1 and, as a matter of consistency, we see the inclusions

$$E_7 \supset E_6 \supset SO(10) \supset SU(5) \supset SU(4) \supset SU(3) \supset SU(2) \supset SU(1),$$

Table 1. The bottoms and related enhanced gauge groups.

<i>H</i>	Bottom
<i>SU</i> (1)	{ <i>pt</i> ₁ ⁽¹⁾ }
<i>SU</i> (2)	{ <i>pt</i> ₁ ⁽¹⁾ , <i>pt</i> ₂ ⁽¹⁾ }
<i>SU</i> (3)	{ <i>pt</i> ₁ ⁽¹⁾ , <i>pt</i> ₂ ⁽¹⁾ , <i>pt</i> ₃ ⁽¹⁾ }
<i>SU</i> (4)	{ <i>pt</i> ₁ ⁽¹⁾ , <i>pt</i> ₂ ⁽¹⁾ , <i>pt</i> ₃ ⁽¹⁾ , <i>pt</i> ₄ ⁽¹⁾ }
<i>SU</i> (5)	{ <i>pt</i> ₁ ⁽¹⁾ , <i>pt</i> ₂ ⁽¹⁾ , <i>pt</i> ₃ ⁽¹⁾ , <i>pt</i> ₄ ⁽¹⁾ , <i>pt</i> ₅ ⁽¹⁾ }
<i>SU</i> (6)	{ <i>pt</i> ₁ ⁽¹⁾ , <i>pt</i> ₂ ⁽¹⁾ , <i>pt</i> ₃ ⁽¹⁾ , <i>pt</i> ₄ ⁽¹⁾ , <i>pt</i> ₅ ⁽¹⁾ , <i>pt</i> ₆ ⁽¹⁾ }
<i>SO</i> (10)	{ <i>pt</i> ₁ ⁽²⁾ , <i>pt</i> ₂ ⁽²⁾ , <i>pt</i> ₃ ⁽¹⁾ , <i>pt</i> ₄ ⁽¹⁾ , <i>pt</i> ₅ ⁽¹⁾ }
<i>E</i> ₆	{ <i>pt</i> ₁ ⁽³⁾ , <i>pt</i> ₂ ⁽²⁾ , <i>pt</i> ₃ ⁽²⁾ , <i>pt</i> ₄ ⁽¹⁾ , <i>pt</i> ₅ ⁽¹⁾ }
<i>E</i> ₇	{ <i>pt</i> ₁ ⁽⁴⁾ , <i>pt</i> ₂ ⁽³⁾ , <i>pt</i> ₃ ⁽²⁾ , <i>pt</i> ₄ ⁽²⁾ , <i>pt</i> ₅ ⁽¹⁾ }
<i>E</i> ₈	{ <i>pt</i> ₁ ⁽⁶⁾ , <i>pt</i> ₂ ⁽⁴⁾ , <i>pt</i> ₃ ⁽³⁾ , <i>pt</i> ₄ ⁽²⁾ , <i>pt</i> ₅ ⁽¹⁾ }

as inclusions of the respective polyhedra. Thus we can see the duality between compactifications of the $E_8 \times E_8$ heterotic string and the type IIA string compactified on a Calabi-Yau manifold. The new contribution here is the observation that sequences of reflexive polyhedra associated to these spaces are nested in such a way as to reflect heterotic perturbative and non-perturbative processes. This qualitative observation is supported by quantitative agreement of the computed from the polyhedra Hodge numbers and the number of tensor, vector and hypermultiplets in the heterotic side. It leads also to the observation that the Dynkin diagrams for the groups may be read off from the polyhedra. Table 2 for the chains of fourfolds demonstrates the nesting of polyhedra.

Table 2. Matter content for the chain of enhanced gauge groups.

Group	Matter content
<i>SU</i> (2)	$(6n + 2)\mathbf{2}$
<i>SU</i> (3)	$(6n + 18)\mathbf{3}$
<i>SU</i> (4)	$(n + 2)\mathbf{6} + (4n + 16)\mathbf{4}$
<i>SU</i> (5)	$(3n + 16)\mathbf{5} + (2 + n)\mathbf{10}$
<i>SO</i> (10)	$(n + 4)\mathbf{16} + (n + 6)\mathbf{10}$
<i>E</i> ₆	$(n + 6)\mathbf{27}$
<i>E</i> ₇	$(\frac{n}{2} + 4)\mathbf{56}$

For the chain $H = SU(1), \dots, SU(5), SO(10), E_6, E_7$ we can determine the number of multiplets of solitonic states that appear from singularities of elliptic fibration. Nesting of polyhedra is interpreted as phase transition. This phase transition can be interpreted as a group chain of spontaneous symmetry breaking

$$SO(10) \rightarrow SU(5) \rightarrow SU(3) \times SU(2) \times U(1)$$

Group $SO(10)$ has group $SU(5)$ as its subgroup:

$$16 = 10 + \bar{5} + 1 .$$

Thus 16 multiplet of fermions consists of three $SU(5)$ multiplets. Particle content of Minimal Supersymmetric Standard Model (MSSM) can be presented in terms of $SU(5)$ multiplets

$$3 \times (24 + 5_H + \bar{5}_H + 5_M + \bar{5}_M + 10_M + \bar{10}_M) ,$$

where 5_H and $\bar{5}_H$ are Higgs multiplets, $\bar{5}_M$ and 10_M are multiplets of quark and lepton superpartners. The transition between $SU(5)$ and $SU(3) \times SU(2) \times U(1)$ can be realised with the help of partial widths of superpartners calculations. Using this restricted parameter set of MSSM model

$$M_0 = 800 \text{ GeV} , \quad M_{1/2} = 650 \text{ GeV} , \\ A_0 = 0 , \quad \tan\beta = 10 , \quad \text{sgn}(\mu) = +1 .$$

it is possible to calculate partial widths of superpartners by application of the computer program SDECAY [12]. These partial widths are shown in Tables 3, 4.

Table 3. Partial widths of superpartners

	channel	BR	channel	BR
\tilde{u}_R	$\tilde{\chi}_1^0 u$	0.997	$\tilde{\chi}_4^0 u$	0.002
\tilde{d}_R	$\tilde{\chi}_1^0 d$	0.997	$\tilde{\chi}_4^0 d$	0.002
\tilde{c}_R	$\tilde{\chi}_1^0 c$	0.997	$\tilde{\chi}_4^0 c$	0.002
\tilde{s}_R	$\tilde{\chi}_1^0 s$	0.997	$\tilde{\chi}_4^0 s$	0.002

Table 4. Partial widths of superpartners

	channel	BR	channel	BR
g	$b_1 b^*$	0.074	$t_1 t^*$	0.425
	$b_1^* b$	0.074	$t_1^* t$	0.425

4 Conclusions

Thus we see the duality between the compactifications of the $E_8 \times E_8$ heterotic string and the type IIA string compactification on a Calabi-Yau manifold. The observations concerning this duality are the subject of the present article: we have a correspondence between reflexive polyhedra, Δ , and Calabi-Yau manifolds. Combining this with the correspondence between vector bundles and Calabi-Yau manifolds gives a correspondence between vector bundles and reflexive polyhedra Δ . It is known also that the moduli spaces of Calabi-Yau manifolds form a web in which continuous transitions between different Calabi-Yau manifolds (phases) occur [13]. Indeed it seems likely that all Calabi-Yau manifolds are connected by processes of this type [14]. In virtue of duality this web structure must exist also on the heterotic side, though the physical picture is different. Indeed, the space of heterotic vacua also forms a web in which different models are connected along branches parametrized by vacuum expectation values of scalars in vector and hypermultiplets. Subsequently, it was argued that the appearance of perturbative heterotic groups in the chains could be explained in the Calabi-Yau picture [6]. Indeed, there is a natural sequence of E-group embeddings which give the Standard Model gauge group and matter structure in an elegant manner:

$$E_3 \times U(1) \subset E_4 \subset E_5 \subset E_6 \subset \dots$$

where $E_3 = SU(3) \times SU(2)$ denotes the non-abelian gauge group of the Standard Model, $E_4 = SU(5)$ and $E_5 = SO(10)$. Both gauge coupling unification as well as the matter content of the Standard Model hint at the presence of a unified gauge group structure at high energies. From this perspective, we interpret the unification of the gauge couplings in the minimal supersymmetric extension of the Standard Model as evidence that instead of the infinitely complex variety of classical groups and matter which can appear in string theory, the list of relevant simple gauge groups is limited to the finite number of exceptional gauge groups and their subgroups. This provides an economical way to break the exceptional gauge group to gauge symmetries closer to the MSSM. One of the nicest qualities of supersymmetry is that so much is known about it already, despite the present lack of direct experimental evidence. The interactions of the Standard Model particles and their superpartners are fixed by supersymmetry, up to mass mixing effects due to supersymmetry breaking. Even the terms and stakes of many of the important outstanding questions, especially the paramount issue "How is supersymmetry broken?", are already rather clear. Analyses of LHC searches corresponding to $\sim 5 \text{ fb}^{-1}$ at $\sqrt{s} = 7 \text{ TeV}$ have not found any superpartners, and have put strong lower bounds on the masses of squarks and gluinos in large classes of models. The most common templates used for reporting the results of experimental searches are the MSUGRA scenario with new parameters $m_0, m_{1/2}, A_0, \tan\beta, \text{Arg}(\mu)$. However, one should not lose sight of the fact that the only indispensable idea of supersymmetry is simply that of a symmetry between fermions and bosons. If supersymmetry is experimentally verified, the discovery will not be an end, but rather a beginning in high energy physics. It seems likely to present us with questions and challenges that we can only guess at presently. The measurement of sparticle masses, production cross-sections, and decay modes will rule out some models for supersymmetry breaking and lend credence to others. We will be able to test the principle of R-parity conservation, the idea that supersymmetry has something to do with the dark matter, and possibly make connections to other aspects of cosmology including baryogenesis and inflation.

REFERENCES

- [1] A.J. Buras et al. Aspects of the grand unification of strong, weak and electromagnetic interactions, Nucl. Phys., Vol. B135, 66–92, 1978.
- [2] M. Kaku. Introduction to Superstring and M-Theory, New York, USA: Springer-Verlag, 1999, 593p.
- [3] C. Vafa. Evidence for F-theory, arXiv: hep-th/9602022.

- [4] N. Arkani-Hamed, S. Dimopoulos and G. Dvali. The Hierarchy problem and new dimensions at a millimeter, *Phys. Lett.*, Vol. B429, 263-272, 1998.
- [5] C. Beasley, Jonathan J. Heckman and C. Vafa. GUTs and Exceptional Branes in F-theory - I , arXiv:0802.3391 [hep-th].
- [6] P. Candelas and A. Font. Duality Between the Webs of Heterotic and Type II Vacua, arXiv: hep-th/9603170.
- [7] E. Perevalov and H. Skarke. Enhanced Gauge Symmetry in Type II and F-Theory Compactifications: Dynkin Diagrams From Polyhedra, arXiv: hep-th/9704129.
- [8] D. R. Morrison and C. Vafa. Compactifications of F-theory on Calabi-Yau threefolds (I), *Nucl. Phys.*, Vol. B473, 74-92, 1996.
- [9] D. R. Morrison and C. Vafa. Compactifications of F-theory on Calabi-Yau threefolds (II), *Nucl. Phys.*, Vol. B476, 437-469, 1996.
- [10] M. Kreuzer and H. Skarke. No Mirror Symmetry in Landau-Ginzburg Spectra!, *Nucl. Phys.*, Vol. B388, 113-130, 1992.
- [11] V. Batyrev. Variations of the mixed Hodge structure of affine hypersurfaces in algebraic tori, *Duke Math. Journ.*, Vol. 69, 349-409, 1993.
- [12] M. Muhlleitner, A. Djouadi and Y. Mambrini. SDECAY: a fortran code for the decays of the supersymmetric particles in the MSSM, *Comput. Phys. Commun.*, Vol. 168, 46-70, 2005.
- [13] M. Reid. The moduli space of 3-folds with $K = 0$ may nevertheless be irreducible, *Math. Ann.*, Vol. 278, 329-334, 1987.
- [14] P. Berglund, S. Katz and A. Klemm. Mirror Symmetry and the Moduli Space for Generic Hypersurfaces in Toric Varieties, *Nucl.Phys.*, Vol. B456, 153-204, 1995.