

# A Joint Chance Constrained Programming with Bivariate Dagum Distribution

Khalid M. El-khabeary\*, Afaf El-Dash, Nada M. Hafez, Samah M. Abo-El-hadid

Department of Mathematics, Insurance and Applied Statistics, Helwan University, Egypt

Received March 5, 2023; Revised August 2, 2023; Accepted August 23, 2023

## Cite This Paper in the Following Citation Styles

(a): [1] Khalid M. El-khabeary, Afaf El-Dash, Nada M. Hafez, Samah M. Abo-El-hadid, "A Joint Chance Constrained Programming with Bivariate Dagum Distribution," *Mathematics and Statistics*, Vol. 11, No. 5, pp. 778 - 785, 2023. DOI: 10.13189/ms.2023.110503.

(b): Khalid M. El-khabeary, Afaf El-Dash, Nada M. Hafez, Samah M. Abo-El-hadid (2023). A Joint Chance Constrained Programming with Bivariate Dagum Distribution. *Mathematics and Statistics*, 11(5), 778 - 785. DOI: 10.13189/ms.2023.110503.

Copyright©2023 by authors, all rights reserved. Authors agree that this article remains permanently open access under the terms of the Creative Commons Attribution License 4.0 International License

**Abstract** A Joint chance-constrained programming (JCCP) technique is regarded as one of the most useful applicable techniques of stochastic programming techniques. It is more suitable for solving uncertain real problems, especially in economics and social problems, where some of model parameters are positive dependent random variables and follow well-known probability distributions. In this paper, we take into account a linear JCCP problem where some right-hand side random parameters are dependent and follow the Dagum distributions. So, firstly we derive a bivariate Dagum distribution with seven parameters with marginals following the Dagum distribution with three parameters. This proposed bivariate Dagum distribution is based on the Farlie-Gumbel-Morgensten copula (as presented in theorem (2.1)). Secondly, the proposed bivariate distribution is used in the context of JCCP technique to transform a linear JCCP model into an exact equivalent deterministic nonlinear programming model through theorem (3.1). Thirdly, through theorem (3.2), we prove that the obtained exact equivalent deterministic nonlinear programming model is a convex model, hence any nonlinear programming method can be used to solve it and find the global optimal solution. Finally, in order to demonstrate how to convert a linear JCCP model into an equivalent deterministic nonlinear programming model and solve it using the cutting plane method, a numerical example is included.

Chance Constrained Programming (CCP), JCCP, Bivariate Dagum Distribution, Multivariate Distributions, Equivalent Deterministic Model, Convex Model, Cutting Plane Method, Global Solution

## 1. Introduction

In 1977, Camilo Dagum first presented the three-parameter Dagum distribution for modelling personal income data [1]. The Pareto and Log-normal distributions can also be replaced with it [2, 3]. Additionally, several applications have used this distribution, for example, income and consumption data [4], financial data [5], and reliability analysis [6]. It is worth noting that the Dagum distribution is related to many distributions, for instance, the Burr type XII, Burr type III, Lomax, log logistic distribution, and the exponential distribution [2,7].

A continuous random variable is said to have a Dagum distribution, or  $X \sim Dag(\beta; \lambda; \delta)$ , if its probability density function (PDF), cumulative function (CDF), and the inverse cumulative function are respectively as follows [1,3]:

$$f(x; \beta; \lambda; \delta) = \beta \lambda \delta x^{-\delta-1} (1 + \lambda x^{-\delta})^{-\beta-1},$$
$$x > 0, \quad \beta, \lambda, \delta > 0, \quad (1)$$

$$F(x; \beta; \lambda; \delta) = (1 + \lambda x^{-\delta})^{-\beta}, \quad (2)$$

**Keywords** Farlie-Gumbel-Morgensten Copula,

$$F^{-1}(y; \beta; \lambda; \delta) = \lambda^{\frac{1}{\delta}} \left( y^{-\frac{1}{\beta}} - 1 \right)^{-\frac{1}{\delta}}, \quad 0 \leq y \leq 1, \tag{3}$$

where  $\beta$  and  $\delta$  are the shape parameters,  $\lambda$  is the scale parameter, and  $y$  is the value of the cumulative function. Now, consider individual chance constraints having only one random parameter on the right-hand side (RHS), as follows:

$$P_r\{\sum_{j=1}^n a_{ij}x_j \geq \tilde{b}_i\} \geq \gamma_i, \quad i = 1, \dots, m_1, \tag{4}$$

$$P_r\{\sum_{j=1}^n a_{ij}x_j \leq \tilde{b}_i\} \geq \gamma_i, \quad i = m_1 + 1, \dots, m_2. \tag{5}$$

Here  $x_j, j = 1, 2, \dots, n$ , represents the decision variable,  $a_{ij}, i = 1, 2, \dots, m_2, j = 1, 2, \dots, n$ , is constant. and  $\tilde{b}_i \sim \text{Dag}(\beta_i, \lambda_i, \delta_i), i = 1, \dots, m_2$ , Finally,  $0 \leq \gamma_i \leq 1$  represents the tolerance measure for the  $i^{th}$  constraint, then the exact equivalent deterministic linear constraints respectively are as follows [8]:

$$\sum_{j=1}^n a_{ij}x_j \geq \lambda_i^{\frac{1}{\delta_i}} \left( \gamma_i^{-\frac{1}{\beta_i}} - 1 \right)^{-\frac{1}{\delta_i}}, \quad i = 1, \dots, m_1, \tag{6}$$

$$\sum_{j=1}^n a_{ij}x_j \leq \lambda_i^{\frac{1}{\delta_i}} \left( (1 - \gamma_i)^{-\frac{1}{\beta_i}} - 1 \right)^{-\frac{1}{\delta_i}}, \quad i = m_1 + 1, \dots, m_2. \tag{7}$$

Recently, many bivariate Dagum distributions were presented. For example, in 2017, Muhammed [9] proposed a singular bivariate Dagum distribution using an idea proposed by Marshall and Olkin. In 2018, Popović et al [10] introduced a tractable bivariate extension of the univariate Dagum distribution by using the Marshall–Olkin approach and examining its dependence properties.

Historically the chance-constrained programming (CCP) technique was introduced by Charnas and Cooper [11] in 1959, in the case of individual chance constraints. Then, in 1965, Miller and Wagner [12] presented the joint chance constraints case. So, since the emergence of this technique, many studies have been presented, and most of those studies assumed the independence between random parameters for simplicity. However, usually many random parameters are dependent and follow non-negative distributions in real-world applications, especially in economic and social problems [8,13].

Therefore, many studies presented the equivalent deterministic programming models corresponding to the JCCP models under the dependence assumption. For example, in 2018, Hafez et al [14,15] assumed that the RHS or LHS parameters were dependent random variables (RVs) that followed the approximated Downton bivariate exponential distribution. In 2019, El-Dash [16] presented an extension of the Freund bivariate exponential distribution, assuming that the parameters in the RHS or some of the LHS parameters are RVs. Then, in 2020, El-Dash presented an extension of Farlie, Gumbel, and Morgenstern's bivariate exponential distribution [17]. In 2021, Ebaid and El-Dash [18] introduced a bivariate generalized exponential by the Cuadras-Augé copula function for the univariate generalized exponential distributions, assuming that the random parameters in the

R.H.S. of joint constraints follow the proposed distribution. Then, in the same year, they assumed that the RHS parameters were dependent RVs that follow a derived bivariate Lomax distribution using the Farlie-Gumbel-Morgenstern (FGM) copula function [19].

The structure of this study is as follows: we present a new bivariate Dagum distribution using FGM Copula in Section 2. Following that, the JCCP linear model with dependent random parameters and its exact equivalent deterministic nonlinear programming model are provided in Section 3. We present a numerical example in Section 4. Finally, section 5 presents conclusions.

## 2. A new Bivariate Dagum Distribution Using FGM Copula

Copula functions are tools for modelling multivariate stochastic dependence between RVs and can be defined as functions that link or join multivariate distributions with their one-dimensional marginal functions. It was first introduced in the work of Sklar [20] in 1959 through his theorem. In literature, there are many families of copulas with one or more dependence parameters, for instance, the Farlie-Gumbel-Morgenstern copula, the Gaussian copula, the Archimedean copula, etc., and they have different properties that are indispensable in many applications [21]. One of the most important families is the FGM copula, which was introduced first by Morgenstern [22].

According to the bivariate FGM copula function, the joint CDF (JCDF) and the FGM copula density function for two RVs  $X_1, X_2$  respectively, are defined by [16]:

$$F(x_1, x_2) = C(F(x_1), G(x_2)) = F(x_1)G(x_2)[1 + \theta(1 - F(x_1))(1 - G(x_2))], \quad \theta \in [-1, 1], \quad (8)$$

$$c(F(x_1), G(x_2)) = [1 + \theta(1 - 2F(x_1))(1 - 2G(x_2))]. \quad (9)$$

Where:  $\theta$  is the dependent parameter which shows the level of dependence between two RVs, so that when it is equal to zero, it means that the two variables are independent.

Also, the joint PDF (JPDF) is as follows:

$$f(x_1, x_2) = f(x_1)g(x_2) \cdot c(F(x_1), G(x_2)). \quad (10)$$

Through the following theorem, the FGM copula will be used to build a new bivariate Dagum distribution, where the JPDF and the JCDF are presented.

**Theorem (2.1):** Let  $X_i \sim \text{Dag}(\beta_i; \lambda_i; \delta_i), i = 1, 2$ , be dependent RVs, then the JCDF and the JPDF of  $(X_1, X_2)$  respectively are as follows:

$$F(x_1, x_2) = \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} + \theta \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} \left\{ 1 - \sum_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} + \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} \right\}, \quad (11)$$

$$f(x_1, x_2) = \prod_{i=1}^2 \beta_i \lambda_i \delta_i x_i^{-\delta_i - 1} (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i - 1} \left[ 1 + \theta \left( 1 - 2 \sum_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} + 4 \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} \right) \right], \quad (12)$$

$$x_i > 0, \quad \beta_i, \lambda_i, \delta_i > 0, \quad i = 1, 2, \quad -1 \leq \theta \leq 1.$$

**Proof:** According to the bivariate FGM copula function, the JCDF is obtained as follows:

$$\begin{aligned} F(x_1, x_2) &= F(x_1)G(x_2)[1 + \theta(1 - F(x_1))(1 - G(x_2))] \\ &= (1 + \lambda_1 x_1^{-\delta_1})^{-\beta_1} (1 + \lambda_2 x_2^{-\delta_2})^{-\beta_2} \left\{ 1 + \theta \left[ 1 - (1 + \lambda_1 x_1^{-\delta_1})^{-\beta_1} \right] \cdot \left[ 1 - (1 + \lambda_2 x_2^{-\delta_2})^{-\beta_2} \right] \right\} \\ &= \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} \left\{ 1 + \theta - \theta \sum_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} + \theta \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} \right\} \\ &= \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} + \theta \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} \left\{ 1 - \sum_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} + \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} \right\}. \end{aligned}$$

Then, the JPDF is obtained as follows:

$$\begin{aligned} f(x_1, x_2) &= f(x_1)g(x_2) \cdot c(F(x_1), G(x_2)) \\ &= \beta_1 \lambda_1 \delta_1 \beta_2 \lambda_2 \delta_2 \cdot x_1^{-\delta_1 - 1} (1 + \lambda_1 x_1^{-\delta_1})^{-\beta_1 - 1} \cdot x_2^{-\delta_2 - 1} (1 + \lambda_2 x_2^{-\delta_2})^{-\beta_2 - 1} \\ &\quad \cdot \left\{ 1 + \theta \left[ 1 - 2(1 + \lambda_1 x_1^{-\delta_1})^{-\beta_1} \right] \cdot \left[ 1 - 2(1 + \lambda_2 x_2^{-\delta_2})^{-\beta_2} \right] \right\} \\ &= \prod_{i=1}^2 \beta_i \lambda_i \delta_i x_i^{-\delta_i - 1} (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i - 1} \cdot \left[ 1 + \theta \left( 1 - 2 \sum_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} + 4 \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} \right) \right]. \end{aligned}$$

The surface plots of the bivariate PDF and CDF of the Dagum distribution (BDagum) for various parameter values  $(\beta_1, \lambda_1, \delta_1, \beta_2, \lambda_2, \delta_2; \theta)$  are shown in Figures 1 and 2, respectively, as follows.




Figure 1. The JPDF plot of the BDagum for various values of parameters  $(\beta_1, \lambda_1, \delta_1, \beta_2, \lambda_2, \delta_2; \theta)$ .




Figure 2. The JCDF plot of the BDagum for various values of parameters  $(\beta_1, \lambda_1, \delta_1, \beta_2, \lambda_2, \delta_2; \theta)$ .

**Special Case (2.1):** In case of independence between RVs  $x_i, i = 1, 2$ , we have,  $\theta = 0$ , and then the JCDF and the JPDF in (11) and (12), respectively, become as follows:

$$F(x_1, x_2) = \prod_{i=1}^2 (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i} \quad (13)$$

$$f(x_1, x_2) = \prod_{i=1}^2 \beta_i \lambda_i \delta_i x_i^{-\delta_i-1} (1 + \lambda_i x_i^{-\delta_i})^{-\beta_i-1}, x_i > 0, \quad \beta_i, \lambda_i, \delta_i > 0, \quad i = 1, 2. \quad (14)$$

In the next section, we will apply the previous theorem in the context of probabilistic programming model.

### 3. Joint Chance Constrained Linear Programming Model

Here, we will illustrate the conversion of the linear JCCP model into an exact equivalent deterministic nonlinear model assuming that some parameters follow the derived bivariate Dagum distribution [8,13].

Consider the following linear JCCP model:

$$Min. Z = \sum_{j=1}^n C_j x_j, \quad (15)$$

$$S. T. \sum_{j=1}^n a_{ij} x_j \geq b_i, \quad i = 1, 2, \dots, m, \quad (16)$$

$$P_r \{ \sum_{j=1}^n a_{ij} x_j \geq \tilde{b}_i, \sum_{j=1}^n a_{i+1,j} x_j \geq \tilde{b}_{i+1} \} \geq \gamma_i, \quad i = m + 1, \quad (17)$$

$$Pr\left\{\sum_{j=1}^n a_{ij}x_j \leq \tilde{b}_i, \sum_{j=1}^n a_{i+1,j}x_j \leq \tilde{b}_{i+1}\right\} \geq \gamma_i, \quad i = m + 2, \tag{18}$$

$$x_j \geq 0, \quad j = 1, \dots, n, \tag{19}$$

where  $C_j, j = 1, 2, \dots, n$ , and  $b_i, i = 1, 2, \dots, m$ , are constants, and  $(\tilde{b}_i, \tilde{b}_{i+1}) \sim BDag(\beta_i, \lambda_i, \delta_i, \beta_{i+1}, \lambda_{i+1}, \delta_{i+1}; \theta), i = m + 1, m + 2$ , are dependent random parameters and follow the above bivariate distribution in (11) and (12).

In the following theorem, we introduce the exact nonlinear equivalent deterministic constraints of the joint chance constraints in (17) and (18).

**Theorem (3.1):** Consider the joint chance constraints in (17) and (18). Then, the exact equivalent deterministic nonlinear constraints respectively are as follows:

$$\prod_{s=i}^{i+1} f_s(x_s)^{-\beta_s} + \theta \prod_{s=i}^{i+1} f_s(x_s)^{-\beta_s} \{1 - \sum_{s=i}^{i+1} f_s(x_s)^{-\beta_s} + \prod_{s=i}^{i+1} f_s(x_s)^{-\beta_s}\} \geq \gamma_i, \quad i = m + 1, \tag{20}$$

$$\prod_{s=i}^{i+1} f_s(x_s)^{-\beta_s} + \theta \prod_{s=i}^{i+1} f_s(x_s)^{-\beta_s} \{1 - \sum_{s=i}^{i+1} f_s(x_s)^{-\beta_s} + \prod_{s=i}^{i+1} f_s(x_s)^{-\beta_s}\} \leq (1 - \gamma_i), \quad i = m + 2, \tag{21}$$

Where  $f_s(x_s) = (1 + \lambda_s(\sum_{j=1}^n a_{sj}x_j)^{-\delta_s})$ .

**Proof:** For the joint chance constraints in (17), it is clear that [8]:

$$Pr\left\{\sum_{j=1}^n a_{ij}x_j \geq \tilde{b}_i, \sum_{j=1}^n a_{i+1,j}x_j \geq \tilde{b}_{i+1}\right\} \geq \gamma_i \rightarrow F_{\tilde{b}_i, \tilde{b}_{i+1}}\left(\sum_{j=1}^n a_{ij}x_j, \sum_{j=1}^n a_{i+1,j}x_j\right) \geq \gamma_i$$

where  $F_{\tilde{b}_i, \tilde{b}_{i+1}}(\dots)$  represents the JCDF of  $\tilde{b}_i, \tilde{b}_{i+1}$  defined in (11). Therefore, by direct substitution, we get constraint (20).

Similarly, for the joint chance constraints in (18) we have,

$$Pr\left\{\sum_{j=1}^n a_{ij}x_j \leq \tilde{b}_i, \sum_{j=1}^n a_{i+1,j}x_j \leq \tilde{b}_{i+1}\right\} \geq \gamma_i \rightarrow F_{\tilde{b}_i, \tilde{b}_{i+1}}\left(\sum_{j=1}^n a_{ij}x_j, \sum_{j=1}^n a_{i+1,j}x_j\right) \leq (1 - \gamma_i),$$

Therefore, by substituting  $F_{\tilde{b}_i, \tilde{b}_{i+1}}(\dots)$ , we get constraint (21).

**Special case (3.1):** When  $\theta = 0$ , then the corresponding exact equivalent deterministic linear constraints of the constraints in (20) and (21) respectively are as follows:

$$0 \leq \sum_{j=1}^n a_{ij}x_j \leq A_i, \text{ and } 0 \leq \sum_{j=1}^n a_{i+1,j}x_j \leq A_{i+1}, \tag{22}$$

$$0 \leq \sum_{j=1}^n a_{ij}x_j \leq B_i, \text{ and } 0 \leq \sum_{j=1}^n a_{i+1,j}x_j \leq B_{i+1}, \tag{23}$$

where the constants are  $A_i = (\lambda_i)^{\frac{1}{\delta_i}} \left( e^{\frac{-\ln(\gamma_i)}{\beta_i}} - 1 \right)^{-\frac{1}{\delta_i}}$ ,  $A_{i+1} = (\lambda_{i+1})^{\frac{1}{\delta_{i+1}}} \left( e^{\frac{-\ln(\gamma_{i+1})}{\beta_{i+1}}} - 1 \right)^{-\frac{1}{\delta_{i+1}}}$ ,

$$B_i = (\lambda_i)^{\frac{1}{\delta_i}} \left( e^{\frac{-\ln(1-\gamma_i)}{\beta_i}} - 1 \right)^{-\frac{1}{\delta_i}}$$
,  $B_{i+1} = (\lambda_{i+1})^{\frac{1}{\delta_{i+1}}} \left( e^{\frac{-\ln(1-\gamma_{i+1})}{\beta_{i+1}}} - 1 \right)^{-\frac{1}{\delta_{i+1}}}$ .

**Proof:** In the case of independence, then  $\theta = 0$  in constraint (20) and the equivalent deterministic nonlinear constraint is as follows:

$$\left[1 + \lambda_i(\sum_{j=1}^n a_{ij}x_j)^{-\delta_i}\right]^{-\beta_i} \cdot \left[1 + \lambda_{i+1}(\sum_{j=1}^n a_{i+1,j}x_j)^{-\delta_{i+1}}\right]^{-\beta_{i+1}} \geq \gamma_i.$$

The previous constraint is a nonlinear constraint, a logarithmic transformation could be applied as follows:

$$-\beta_i \ln \left[1 + \lambda_i \left(\sum_{j=1}^n a_{ij}x_j\right)^{-\delta_i}\right] - \beta_{i+1} \ln \left[1 + \lambda_{i+1} \left(\sum_{j=1}^n a_{i+1,j}x_j\right)^{-\delta_{i+1}}\right] \geq \ln(\gamma_i).$$

Let  $h_i = \ln \left[1 + \lambda_i (\sum_{j=1}^n a_{ij}x_j)^{-\delta_i}\right]$  and  $h_{i+1} = \ln \left[1 + \lambda_{i+1} (\sum_{j=1}^n a_{i+1,j}x_j)^{-\delta_{i+1}}\right]$  (24)

$$\therefore \beta_i h_i + \beta_{i+1} h_{i+1} \leq -\ln(\gamma_i).$$

Therefore,

$$0 \leq h_i \leq \frac{-\ln(\gamma_i)}{\beta_i} \quad \text{and} \quad 0 \leq h_{i+1} \leq \frac{-\ln(\gamma_{i+1})}{\beta_{i+1}}. \tag{25}$$

Using (24), (25) we get

$$\sum_{j=1}^n a_{ij}x_j = (\lambda_i)^{\frac{1}{\delta_i}}(e^{h_i} - 1)^{-\frac{1}{\delta_i}} = (\lambda_i)^{\frac{1}{\delta_i}} \left( e^{\frac{-\ln(\gamma_i)}{\beta_i}} - 1 \right)^{-\frac{1}{\delta_i}}.$$

Hence

$$0 \leq \sum_{j=1}^n a_{ij}x_j \leq A_i, \tag{26}$$

Also,

$$0 \leq \sum_{j=1}^n a_{i+1,j}x_j \leq A_{i+1}, \tag{27}$$

where  $A_i = (\lambda_i)^{\frac{1}{\delta_i}} \left( e^{\frac{-\ln(\gamma_i)}{\beta_i}} - 1 \right)^{-\frac{1}{\delta_i}}$ ,  $A_{i+1} = (\lambda_{i+1})^{\frac{1}{\delta_{i+1}}} \left( e^{\frac{-\ln(\gamma_{i+1})}{\beta_{i+1}}} - 1 \right)^{-\frac{1}{\delta_{i+1}}}$

Similarly, for constraint (21), the exact equivalent deterministic linear constraint is as follows:

$$0 \leq \sum_{j=1}^n a_{ij}x_j \leq B_i, \tag{28}$$

Also,

$$0 \leq \sum_{j=1}^n a_{i+1,j}x_j \leq B_{i+1} \tag{29}$$

where  $B_i = (\lambda_i)^{\frac{1}{\delta_i}} \left( e^{\frac{-\ln(1-\gamma_i)}{\beta_i}} - 1 \right)^{-\frac{1}{\delta_i}}$ ,  $B_{i+1} = (\lambda_{i+1})^{\frac{1}{\delta_{i+1}}} \left( e^{\frac{-\ln(1-\gamma_{i+1})}{\beta_{i+1}}} - 1 \right)^{-\frac{1}{\delta_{i+1}}}$ .

**Theorem (3.2):** Consider the nonlinear function in the LHS for constraint (20) or (21). Then the exact equivalent deterministic nonlinear programming model in (15)-(16) and (19)-(21) is a convex model.

**Proof:** The model in (15) -(16) and (19) -(21) is a convex model if and only if the LHS of constraints (20) or (21) are strictly convex functions, which occurs if the following condition is met [23]:

$$F[\alpha x^{(1)} + (1 - \alpha)x^{(2)}] < \alpha F(x^{(1)}) + (1 - \alpha)F(x^{(2)})$$

Then let  $x^{(1)} = \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}$ ,  $x^{(2)} = \begin{pmatrix} X_1 + \Delta_1 \\ X_2 + \Delta_2 \end{pmatrix}$ ,  $x^{(1)} \neq x^{(2)}$ ,  $x^{(1)}, x^{(2)} \geq 0$ ,

Now, by substituting the LHS of either constraint (20) or (21)- because both constraints have the same LHS, into the LHS of the previous condition, then,

$$\begin{aligned} \text{LHS} &= F[\alpha x^{(1)} + (1 - \alpha)x^{(2)}] = F\left[\alpha \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} + (1 - \alpha) \begin{pmatrix} X_1 + \Delta_1 \\ X_2 + \Delta_2 \end{pmatrix}\right] = F\left[\begin{pmatrix} \alpha X_1 \\ \alpha X_2 \end{pmatrix} + \begin{pmatrix} (1 - \alpha)(X_1 + \Delta_1) \\ (1 - \alpha)(X_2 + \Delta_2) \end{pmatrix}\right] = F\left(\begin{pmatrix} X_1 - \alpha\Delta_1 + \Delta_1 \\ X_2 - \alpha\Delta_2 + \Delta_2 \end{pmatrix}\right) \\ &= (1 + \theta) \left(1 + \frac{\lambda_1}{(X_1 - \alpha\Delta_1 + \Delta_1)^{\delta_1}}\right)^{-\beta_1} \left(1 + \frac{\lambda_2}{(X_2 - \alpha\Delta_2 + \Delta_2)^{\delta_2}}\right)^{-\beta_2} - \theta \left(1 + \frac{\lambda_1}{(X_1 - \alpha\Delta_1 + \Delta_1)^{\delta_1}}\right)^{-\beta_1} \left(1 + \frac{\lambda_2}{(X_2 - \alpha\Delta_2 + \Delta_2)^{\delta_2}}\right)^{-2\beta_2} - \theta \left(1 + \frac{\lambda_1}{(X_1 - \alpha\Delta_1 + \Delta_1)^{\delta_1}}\right)^{-2\beta_1} \left(1 + \frac{\lambda_2}{(X_2 - \alpha\Delta_2 + \Delta_2)^{\delta_2}}\right)^{-\beta_2} + \theta \left(1 + \frac{\lambda_1}{(X_1 - \alpha\Delta_1 + \Delta_1)^{\delta_1}}\right)^{-2\beta_1} \cdot \left(1 + \frac{\lambda_2}{(X_2 - \alpha\Delta_2 + \Delta_2)^{\delta_2}}\right)^{-2\beta_2}. \end{aligned}$$

Using the binomial series, where  $(1 + x)^\varphi = \sum_{n=0}^\infty C_n^\varphi x^n$ ,  $\varphi$  is real number,  $|x| < 1$ ,

$$\begin{aligned} \text{LHS} &= \left\{ (1 + \theta) \sum_{n=0}^\infty C_n^{-\beta_1} \sum_{m=0}^\infty C_m^{-\beta_2} + \theta \sum_{n=0}^\infty C_n^{-2\beta_1} \sum_{m=0}^\infty C_m^{-2\beta_2} - \theta \sum_{n=0}^\infty C_n^{-\beta_1} \sum_{m=0}^\infty C_m^{-2\beta_2} - \theta \sum_{n=0}^\infty C_n^{-2\beta_1} \sum_{m=0}^\infty C_m^{-\beta_2} \right\} \left\{ \left[ \frac{\lambda_1}{(X_1 - \alpha\Delta_1 + \Delta_1)^{\delta_1}} \right]^n \left[ \frac{\lambda_2}{(X_2 - \alpha\Delta_2 + \Delta_2)^{\delta_2}} \right]^m \right\}, \tag{30} \end{aligned}$$

Similarly, by substituting the LHS of constraint (20) or (21) into the RHS in the convexity condition, then:

$$\begin{aligned}
 \text{RHS} &= \alpha F(x^{(1)}) + (1 - \alpha)F(x^{(2)}) = \alpha f\left(\frac{X_1}{X_2}\right) + (1 - \alpha) f\left(\frac{(X_1 + \Delta_1)}{(X_2 + \Delta_2)}\right) \\
 &= \alpha \left\{ (1 + \theta) \left(1 + \frac{\lambda_1}{X_1 \delta_1}\right)^{-\beta_1} \left(1 + \frac{\lambda_2}{X_2 \delta_2}\right)^{-\beta_2} - \theta \left(1 + \frac{\lambda_1}{X_1}\right)^{-\beta_1} \left(1 + \frac{\lambda_2}{X_2}\right)^{-\beta_2} - \theta \left(1 + \frac{\lambda_1}{X_1 \delta_1}\right)^{-2\beta_1} \left(1 + \frac{\lambda_2}{X_2 \delta_2}\right)^{-\beta_2} \right. \\
 &\quad \left. + \theta \left(1 + \frac{\lambda_1}{X_1 \delta_1}\right)^{-2\beta_1} \left(1 + \frac{\lambda_2}{X_2 \delta_2}\right)^{-2\beta_2} \right\} \\
 &\quad + (1 - \alpha) \left\{ (1 + \theta) \cdot \left(1 + \frac{\lambda_1}{(X_1 + \Delta_1) \delta_1}\right)^{-\beta_1} \left(1 + \frac{\lambda_2}{(X_2 + \Delta_2) \delta_2}\right)^{-\beta_2} - \theta \left(1 + \frac{\lambda_1}{(X_1 + \Delta_1)}\right)^{-\beta_1} \right. \\
 &\quad \cdot \left(1 + \frac{\lambda_2}{(X_2 + \Delta_2)}\right)^{-\beta_2} - \theta \left(1 + \frac{\lambda_1}{(X_1 + \Delta_1) \delta_1}\right)^{-2\beta_1} \left(1 + \frac{\lambda_2}{(X_2 + \Delta_2) \delta_2}\right)^{-\beta_2} \\
 &\quad \left. + \theta \left(1 + \frac{\lambda_1}{(X_1 + \Delta_1) \delta_1}\right)^{-2\beta_1} \left(1 + \frac{\lambda_2}{(X_2 + \Delta_2) \delta_2}\right)^{-2\beta_2} \right\}.
 \end{aligned}$$

Similarly using the binomial series, we get

$$\begin{aligned}
 \text{RHS} &= \left\{ (1 + \theta) \sum_{n=0}^{\infty} C_n^{-\beta_1} \sum_{m=0}^{\infty} C_m^{-\beta_2} + \theta \sum_{n=0}^{\infty} C_n^{-2\beta_1} \sum_{m=0}^{\infty} C_m^{-2\beta_2} - \theta \sum_{n=0}^{\infty} C_n^{-\beta_1} \cdot \sum_{m=0}^{\infty} C_m^{-2\beta_2} - \right. \\
 &\quad \left. \theta \sum_{n=0}^{\infty} C_n^{-2\beta_1} \sum_{m=0}^{\infty} C_m^{-\beta_2} \right\} \left\{ \alpha \left[ \frac{\lambda_1}{X_1 \delta_1} \right]^n \left[ \frac{\lambda_2}{X_2 \delta_2} \right]^m + \left[ \frac{\lambda_1}{(X_1 + \Delta_1) \delta_1} \right]^n \left[ \frac{\lambda_2}{(X_2 + \Delta_2) \delta_2} \right]^m - \alpha \left[ \frac{\lambda_1}{(X_1 + \Delta_1) \delta_1} \right]^n \left[ \frac{\lambda_2}{(X_2 + \Delta_2) \delta_2} \right]^m \right\}. \quad (31)
 \end{aligned}$$

From (30) and (31), it is clear that the condition is satisfied.

Thus, the function on the LHS of the constraint (20) or (21) is a strictly convex function. So, the model in (15)-(16), (19)-(21) is a strictly convex model. Also, this theorem satisfies the case of independence in the previous special case. This theorem is important because the convex model can be solved using the cutting plane method and get the global optimal solution [23,24].

### 4. A Numerical Example

Consider the following JCCP model:

$$\text{Min. } Z = 2x_1 + x_2, \tag{32}$$

$$\text{S. T. } 2x_1 + x_2 \geq 6, \tag{33}$$

$$x_1 + x_2 \leq 5, \tag{34}$$

$$P_r\{4x_1 + 2x_2 \geq \widetilde{b}_3, 2x_1 + x_2 \geq \widetilde{b}_4\} \geq 0.9, \tag{35}$$

$$x_j \geq 0, j = 1,2, \tag{36}$$

where  $(\widetilde{b}_3, \widetilde{b}_4) \sim BDag(\beta_1 = 1, \lambda_1 = 2, \delta_1 = 2, \beta_2 = 2, \lambda_2 = 1, \delta_2 = 2, \theta = 0.8)$  are dependent random parameters and follow the derived bivariate Dagum distribution.

According to theorem (3.1), the exact equivalent deterministic nonlinear programming model using the JCCP technique, which corresponds to the above model, is as follows:

$$\text{Min. } Z = 2x_1 + x_2, \tag{37}$$

$$\text{S. T. } 2x_1 + x_2 \geq 6, \tag{38}$$

$$x_1 + x_2 \leq 5, \tag{39}$$

$$\frac{1.8(1+2(4x_1+2x_2)^{-2})^{-1}}{(1+(2x_1+x_2)^{-2})^2} - \frac{0.8(1+2(4x_1+2x_2)^{-2})^{-1}}{(1+(2x_1+x_2)^{-2})^4} - \frac{0.8(1+2(4x_1+2x_2)^{-2})^{-2}}{(1+(2x_1+x_2)^{-2})^2} + \frac{0.8(1+2(4x_1+2x_2)^{-2})^{-2}}{(1+(2x_1+x_2)^{-2})^4} \geq 0.9, \tag{40}$$

The above model is convex according to theorem (3.2), therefore it can be solved by the cutting plane method [12,22], assuming that the bounds of the decision variables are

$$0.1 \leq x_1 \leq 4, 0.1 \leq x_2 \leq 3 \tag{41}$$

Then, the global optimal solution is as follows:

$$x_1 = 2.95, x_2 = 0.1, z^* = 6$$

## 5. Conclusions

Throughout this paper, the JPfD and the JCDF of the new bivariate Dagum distribution are presented through theorem (2.1) by using the FGM copula. Then, the JCCP technique is used to transform JCCP model into an exact nonlinear equivalent deterministic programming model through theorem (3.1), where some parameters in the JCCP linear model follow the derived bivariate Dagum distribution, then a special case is introduced regarding independent parameters. After that, we have proved that the obtained exact nonlinear equivalent deterministic programming model is a convex model through theorem (3.2) and a global optimal solution could be obtained. Finally, a numerical example is presented.

---

## REFERENCES

- [1] Dagum, C., "A new model of personal income distribution: specification and estimation. In Modeling income distributions and Lorenz curves," (3-25), New York, NY: Springer New York, 2008.
- [2] Chotikapanich, D. (Ed.), "Modeling income distributions and Lorenz curves (Vol. 5)," Springer Science & Business Media, 2008.
- [3] Dey, S., Al-Zahrani, B., & Basloom, S. "Dagum distribution: Properties and different methods of estimation," *International Journal of Statistics and Probability*, 6(2), 74-92, 2017.
- [4] Domma, F., & Giordano, S., "A stress-strength model with dependent variables to measure household financial fragility," *Statistical Methods & Applications*, 21, 375-389, 2012.
- [5] Ishaq, A. I., & Abiodun, A. A., "A new generalization of Dagum distribution with application to financial data sets. In 2020 International Conference on Data Analytics for Business and Industry: Way Towards a Sustainable Economy (ICDABI)," (1-6), IEEE, 2020.
- [6] Rathie, P. N., de Sena Monteiro Ozelim, L. C., & de Andrade, B. B., "Portfolio Management of Copula-Dependent Assets Based on P (Y < X) Reliability Models: Revisiting Frank Copula and Dagum Distributions," *Stats*, 4(4), 1027-1050, 2021.
- [7] Tadikamalla, P. R., "A look at the Burr and related distributions," *International Statistical Review/Revue Internationale de Statistique*, 337-344, 1980.
- [8] Prekopa, A., "Stochastic Programming," Kluwer Academic Publishers. Dordrecht, The Netherlands, 1995.
- [9] Muhammed, H. Z., "Bivariate dagum distribution," *International Journal of Reliability and Applications*, 18(2), 65-82, 2017.
- [10] Popović, B. V., Genç, A. İ., & Domma, F., "Copula-based properties of the bivariate Dagum distribution," *Computational and Applied Mathematics*, 37, 6230-6251, 2018.
- [11] Charnes, A., & Cooper, W. W., "Chance-constrained programming," *Management science*, 6(1), 73-79, 1959.
- [12] Miller, B. L., & Wagner, H. M., "Chance constrained programming with joint constraints," *Operations Research*, 13(6), 930-945, 1965.
- [13] Birge, J. R., & Louveaux, F., "Introduction to stochastic programming," Springer Science & Business Media, 2011.
- [14] Hafez, N., El-Dash, A., & Albehery, N., "Joint chance constrained programming with dependent parameters," *Advances and Application in Statistics*, 53(1), 1-12, 2018.
- [15] Hafez, N., El-Dash, A., & Albehery, N., "Chance Constrained Programming (CCP) with Independent or Dependent Exponential Input Coefficients," *International Journal of Research and Reviews in Applied Sciences*, 34(2), 2018.
- [16] El-Dash, A., "Probabilistic Linear Problems with Bivariate Exponential Distributed Random Parameters," *International Journal of Research and Reviews in Applied Sciences*, 41(1), 2019.
- [17] El-Dash, A., "Probabilistic programming approach with dependent random parameters," *International Journal of Contemporary Mathematical Sciences*, 15(2), 73-80, 2020.
- [18] Ebaid, R., & El-Dash, A., "A new bivariate generalized exponential distribution with application in joint chance constrained programming," *Far East Journal of Mathematical Sciences*, 130(2), 135-149, 2021.
- [19] Ebaid, R., & El-Dash, A., "a probabilistic programming technique for bivariate lomax random parameters," *Far East Journal of Mathematical Sciences*, 130(2), 151-164, 2021.
- [20] Sklar, M., "Fonctions de répartition à dimensions et leurs marges. In Annales de l'ISUP," 8(3), 229-231, 1959.
- [21] Nelsen, R. B., "An introduction to copulas," Springer & Verla, New York, Inc, 1999.
- [22] Morgenstern, D., "Einfache beispiele zweidimensionaler verteilungen," *Mitteilungsblatt für Mathematische Statistik*, 8, 234-235, 1956.
- [23] Ruszczyński, A., "Nonlinear optimization," Princeton university press, 2011.
- [24] Kelley, Jr, J. E., "The cutting-plane method for solving convex programs," *Journal of the society for Industrial and Applied Mathematics*, 8(4), 703-712, 1960.